



US010376563B1

(12) **United States Patent**
Van Pijkeren et al.

(10) **Patent No.:** **US 10,376,563 B1**
(45) **Date of Patent:** **Aug. 13, 2019**

(54) **METHODS FOR SYSTEMICALLY DELIVERING POLYPEPTIDES AND MICROORGANISMS THEREFOR**

(71) Applicant: **WISCONSIN ALUMNI RESEARCH FOUNDATION**, Madison, WI (US)

(72) Inventors: **Jan Peter Van Pijkeren**, Madison, WI (US); **Alan Attie**, Madison, WI (US); **Mark Keller**, McFarland, WI (US); **Jee-Hwan Oh**, Madison, WI (US)

(73) Assignee: **WISCONSIN ALUMNI RESEARCH FOUNDATION**, Madison, WI (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/431,373**

(22) Filed: **Feb. 13, 2017**

Related U.S. Application Data

(60) Provisional application No. 62/294,578, filed on Feb. 12, 2016.

(51) **Int. Cl.**

C12N 15/74 (2006.01)
A61K 35/74 (2015.01)
A61K 38/00 (2006.01)
A61K 39/09 (2006.01)
A61K 38/20 (2006.01)
A61K 35/747 (2015.01)
A61K 38/02 (2006.01)
A61K 9/00 (2006.01)
C07K 14/54 (2006.01)
A61K 35/00 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 38/20** (2013.01); **A61K 9/0053** (2013.01); **A61K 35/747** (2013.01); **A61K 38/02** (2013.01); **C07K 14/54** (2013.01); **A61K 2035/11** (2013.01)

(58) **Field of Classification Search**

CPC C12N 15/74; C12N 15/746; A61K 35/74; A61K 38/00; A61K 39/09
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,344,867 B2 3/2008 Connolly
9,103,831 B2 8/2015 O'Sullivan et al.
2012/0189550 A1* 7/2012 O'Sullivan A61K 35/74
424/9.6

OTHER PUBLICATIONS

McGee et al., IL-22 Promotes Fibroblast-Mediated Wound Repair in the Skin *Journal of Investigative Dermatology* (2013) 133, 1321-1329.*

Ahrné, S., Molin, G., & Axelsson, L.; Transformation of *Lactobacillus reuteri* with electroporation: Studies on the erythromycin resis-

tance plasmid pLUL631; *Current Microbiology* (1992), 24: 199-205.

Alvarez-Sieiro P, Montalbán-López M, Mu D, Kuipers OP; Bacteriocins of lactic acid bacteria: extending the family. *Appl Microbiol Biotechnol.* 2016. 100(7):2939-51.

Bahey-El-Din, Mohammed, Cormac GM Gahan, and Brendan T. Griffin.; *Lactococcus lactis* as a cell factory for delivery of therapeutic proteins.; *Current gene therapy* 10.1 (2010): 34-45.

Barrangou, Rodolphe, and Jan-Peter van Pijkeren. "Exploiting CRISPR—Cas immune systems for genome editing in bacteria." *Current opinion in biotechnology* 37 (2016): 61-68.

Becker, Stephen C., et al. "LysK CHAP endopeptidase domain is required for lysis of live staphylococcal cells." *FEMS microbiology letters* 294.1 (2009): 52-60.

Beisel, Chase L., Ahmed A. Gomaa, and Rodolphe Barrangou. "A CRISPR design for next-generation antimicrobials." *Genome biology* 15.11 (2014): 516.

Bikard, David, et al. "Exploiting CRISPR—Cas nucleases to produce sequence-specific antimicrobials." *Nature biotechnology* 32.11 (2014): 1146-1150.

Borysowski, Jan, Beata Weber-Dąbrowska, and Andrzej Górski. "Bacteriophage endolysins as a novel class of antibacterial agents." *Experimental Biology and Medicine* 231.4 (2006): 366-377.

Bourguet, Feliza A., et al. "Characterization of a novel lytic protein encoded by the *Bacillus cereus* E33L gene ampD as a *Bacillus anthracis* antimicrobial protein." *Applied and environmental microbiology* 78.8 (2012): 3025-3027.

Britton, Robert A., et al. "Probiotic *L. reuteri* treatment prevents bone loss in a menopausal ovariectomized mouse model." *Journal of cellular physiology* 229.11 (2014): 1822-1830.

Chatel, J. M., et al. "In vivo transfer of plasmid from food-grade transiting lactococci to murine epithelial cells." *Gene therapy* 15.16 (2008): 1184-1190.

Cheng, Xiaodong, et al. "The structure of bacteriophage T7 lysozyme, a zinc amidase and an inhibitor of T7 RNA polymerase." *Proceedings of the National Academy of Sciences* 91.9 (1994): 4034-4038.

Citorik, Robert J., Mark Mimee, and Timothy K. Lu. "Sequence-specific antimicrobials using efficiently delivered RNA-guided nucleases." *Nature biotechnology* 32.11 (2014): 1141-1145.

Cotter, Paul D., R. Paul Ross, Colin Hill. "Bacteriocins—a viable alternative to antibiotics?." *Nature Reviews Microbiology* 11.2 (2013): 95-105.

Cronin, Michelle, et al. "High resolution in vivo bioluminescent imaging for the study of bacterial tumour targeting." *PloS one* 7.1 (2012): e30940.

Cronin, Michelle, et al. "Orally administered bifidobacteria as vehicles for delivery of agents to systemic tumors." *Molecular Therapy* 18.7 (2010): 1397-1407.

(Continued)

Primary Examiner — Maria G Leavitt

(74) *Attorney, Agent, or Firm* — Daniel A. Blasiolo; DeWitt LLP

(57)

ABSTRACT

Methods and microorganisms for systemically introducing a polypeptide in the bloodstream of a subject. The methods of the invention include administering into the gastrointestinal tract of a subject a bacterium configured to express and produce and release the polypeptide. The bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream of the subject, preferably in a detectable amount. The microorganisms of the invention include lactic acid bacteria, such as *Lactobacillus reuteri*, that comprise a recombinant gene configured to express a polypeptide to be systemically introduced.

11 Claims, 12 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

- De Azevedo, Marcela, et al. "In vitro and in vivo characterization of DNA delivery using recombinant *Lactococcus lactis* expressing a mutated form of *L. monocytogenes* Internalin A." *BMC microbiology* 12.1 (2012): 299.
- De Ruyter, P. G., Oscar P. Kuipers, and Willem M. De Vos. "Controlled gene expression systems for *Lactococcus lactis* with the food-grade inducer nisin." *Applied and environmental microbiology* 62.10 (1996): 3662-3667.
- De Weirtdt, Rosemarie, et al. "Glycerol supplementation enhances *L. reuteri*'s protective effect against *S. Typhimurium* colonization in a 3-D model of colonic epithelium." *PLoS One* 7.5 (2012): e37116.
- Dishisha, Tarek, et al. "Flux analysis of the *Lactobacillus reuteri* propanediol-utilization pathway for production of 3-hydroxypropionaldehyde, 3-hydroxypropionic acid and 1, 3-propanediol from glycerol." *Microbial cell factories* 13.1 (2014): 76.
- Doleyres, Y., et al. "Production of 3-hydroxypropionaldehyde using a two-step process with *Lactobacillus reuteri*." *Applied microbiology and biotechnology* 68.4 (2005): 467-474.
- Eaton, Kathryn A., et al. "Probiotic *Lactobacillus reuteri* ameliorates disease due to enterohemorrhagic *Escherichia coli* in germfree mice." *Infection and immunity* 79.1 (2011): 185-191.
- Elzagheid, Adam, et al. "E-cadherin expression pattern in primary colorectal carcinomas and their metastases reflects disease outcome." *World journal of gastroenterology: WJG* 12.27 (2006): 4304.
- Field, Des, et al. "Bioengineered nisin A derivatives with enhanced activity against both Gram positive and Gram negative pathogens." *PLoS One* 7.10 (2012): e46884.
- Field, Des, et al. "The generation of nisin variants with enhanced activity against specific gram-positive pathogens." *Molecular microbiology* 69.1 (2008): 218-230.
- Fischetti, Vincent A. "12 The Use of Phage Lytic Enzymes to Control Bacterial Infections." *Bacteriophages: Biology and applications* (2004): 321.
- Fischetti, Vincent A. "Bacteriophage Lysins: the Ultimate Enzybiotic." *Enzybiotics: Antibiotic Enzymes as Drugs and Therapeutics* (2010): 107-122.
- Fogel, M. R., and Gary M. Gray. "Starch hydrolysis in man: an intraluminal process not requiring membrane digestion." *Journal of applied physiology* 35.2 (1973): 263-267.
- Foster, Patricia L. "Methods for determining spontaneous mutation rates." *Methods in enzymology* 409 (2006): 195-213.
- Frese, Steven A., et al. "The evolution of host specialization in the vertebrate gut symbiont *Lactobacillus reuteri*." *PLoS genetics* 7.2 (2011): e1001314.
- Fu, Geng-Feng, et al. "Bifidobacterium longum as an oral delivery system of endostatin for gene therapy on solid liver cancer." *Cancer gene therapy* 12.2 (2005): 133-140.
- Gibson, Daniel G., et al. "Enzymatic assembly of DNA molecules up to several hundred kilobases." *Nature methods* 6.5 (2009): 343-345.
- Gomaa, Ahmed A., et al. "Programmable removal of bacterial strains by use of genome-targeting CRISPR—Cas systems." *MBio* 5.1 (2014): e00928-13.
- Green et al., *Molecular Cloning: A Laboratory Manual*, 4th ed., Cold Spring Harbor Laboratory Press, 2012.
- Guimarães, Valeria Dellaretti, et al. "Internalin-expressing *Lactococcus lactis* is able to invade small intestine of guinea pigs and deliver DNA into mammalian epithelial cells." *Microbes and infection* 7.5 (2005): 836-844.
- Guimarães, Valeria, et al. "A new plasmid vector for DNA delivery using lactococci." *Genetic vaccines and therapy* 7.1 (2009): 4.
- Hill, C., Guarner, F., Reid, G., Gibson, G.R., Merenstein, D.J., Pot, B., Morelli, L., Canani, R.B., Flint, H.J., Salminen, S. and Calder, P.C. (2014). Expert consensus document: The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nature reviews Gastroenterology & hepatology*, 11(8),506-514.
- Huyghebaert, Nathalie, et al. "Development of an enteric-coated formulation containing freeze-dried, viable recombinant *Lactococcus lactis* for the ileal mucosal delivery of human interleukin-10." *European Journal of Pharmaceutics and Biopharmaceutics* 60.3 (2005): 349-359.
- Jacobs, C., et al. "Bacterial cell wall recycling provides cytosolic muropeptides as effectors for beta-lactamase induction." *The EMBO journal* 13.19 (1994): 4684.
- Jacobs, Christine, et al. "AmpD, essential for both β -lactamase regulation and cell wall recycling, is a novel cytosolic N-acetylmuramyl-L-alanine amidase." *Molecular microbiology* 15.3 (1995): 553-559.
- Jensen, Hanne, et al. "Role of *Lactobacillus reuteri* cell and mucus-binding protein A (CmbA) in adhesion to intestinal epithelial cells and mucus in vitro." *Microbiology* 160.4 (2014): 671-681.
- Kok, Stefan de, et al. "Rapid and reliable DNA assembly via ligase cycling reaction." *ACS synthetic biology* 3.2 (2014): 97-106.
- Kommineni, Sushma, et al. "Bacteriocin production augments niche competition by enterococci in the mammalian gastrointestinal tract." *Nature* 526.7575 (2015): 719-722.
- Lecuit, M. A. R. C., et al. "Internalin of *Listeria monocytogenes* with an intact leucine-rich repeat region is sufficient to promote internalization." *Infection and immunity* 65.12 (1997): 5309-5319.
- Li, C., Chen, X., Kou, L., Hu, B., Zhu, L., Fan, Y. . . Xu, G. (2010); Selenium-Bifidobacterium longum as a delivery system of endostatin for inhibition of pathogenic bacteria and selective regression of solid tumor; *Experimental and Therapeutic Medicine*, 1, 129-135.
- Liu, Yuying, et al. "Human-derived probiotic *Lactobacillus reuteri* strains differentially reduce intestinal inflammation." *American Journal of Physiology-Gastrointestinal and Liver Physiology* 299.5 (2010): G1087-G1096.
- Loessner, Martin J. "Bacteriophage endolysins—current state research and applications." *Current opinion in microbiology* 8.4 (2005): 480-487.
- Loessner, Martin J., et al. "C-terminal domains of *Listeria monocytogenes* bacteriophage murein hydrolases determine specific recognition and high-affinity binding to bacterial cell wall carbohydrates." *Molecular microbiology* 44.2 (2002): 335-349.
- Loonen, L. MP, et al. "REG3 γ -deficient mice have altered mucus distribution and increased mucosal inflammatory responses to the microbiota and enteric pathogens in the ileum." *Mucosal immunology* 7.4 (2014): 939-947.
- Loonen, Linda MP. *RegIII proteins as gatekeepers of the intestinal epithelium*. 2013 Thesis, Wageningen University, 2013. 208 pages.
- López, Rubens, et al. "The pneumococcal cell wall degrading enzymes: a modular design to create new lysins?." *Microbial Drug Resistance* 3.2 (1997): 199-211.
- MacKenzie, Donald A., et al. "Strain-specific diversity of mucus-binding proteins in the adhesion and aggregation properties of *Lactobacillus reuteri*." *Microbiology* 156.11 (2010): 3368-3378.
- Navarre, William Wiley, et al. "Multiple Enzymatic Activities of the Murein Hydrolase from Staphylococcal Phage ϕ 11 Identification of a d-Alanyl-Glycine Endopeptidase Activity." *Journal of Biological Chemistry* 274.22 (1999): 15847-15856.
- Nelson, Daniel, et al. "PlyC: a multimeric bacteriophage lysin." *Proceedings of the National Academy of Sciences* 103.28 (2006): 10765-10770.
- Nicoletti, Mauro, and Giuseppe Elertani. "DNA fusion product of phage P2 with plasmid pBR1022: a new phasmid." *Molecular and General Genetics MGG* 189.2 (1983): 343-347.
- Oh, Jee-Hwan, and Jan-Peter van Pijkeren. "CRISPR—Cas9-assisted recombineering in *Lactobacillus reuteri*." *Nucleic acids research* 42.17 (2014): e131-e131.
- Oh, Phaik Lyn, et al. "Diversification of the gut symbiont *Lactobacillus reuteri* as a result of host-driven evolution." *The ISME journal* 4.3 (2010): 377-387.
- Paez-Espino, David, et al. "Strong bias in the bacterial CRISPR elements that confer immunity to phage." *Nature communications* 4 (2013): 1430.
- Palfy, R., et al. "Bacteria in gene therapy: bactofection versus alternative gene therapy." *Gene Therapy* 13.2 (2006): 101-105.

(56)

References Cited

OTHER PUBLICATIONS

- Pfaffl, Michael W. "A new mathematical model for relative quantification in real-time RT-PCR." *Nucleic acids research* 29.9 (2001): e45-e45.
- Pfaffl, Michael W., Graham W. Horgan, and Leo Dempfle. "Relative expression software tool (REST©) for group-wise comparison and statistical analysis of relative expression results in real-time PCR." *Nucleic acids research* 30.9 (2002): e36-e36.
- Rajpal, Gautam, et al. "Single-chain insulins as receptor agonists." *Molecular endocrinology* 23.5 (2009): 679-688.
- Riboulet-Bisson, Eliette, et al. "Effect of Lactobacillus salivarius bacteriocin Abp118 on the mouse and pig intestinal microbiota." *PLoS one* 7.2 (2012): e31113.
- Riedel, Christian U., et al. "Improved luciferase tagging system for Listeria monocytogenes allows real-time monitoring in vivo and in vitro." *Applied and environmental microbiology* 73.9 (2007): 3091-3094.
- Robert, Sofie, and Lothar Steidler. "Recombinant Lactococcus lactis can make the difference in antigen-specific immune tolerance induction, the Type 1 Diabetes case." *Microbial cell factories* 13.1 (2014): S11.
- Rosche, William A., and Patricia L. Foster. "Determining mutation rates in bacterial populations." *Methods* 20.1 (2000): 4-17.
- Sabat, Robert, Wenjun Ouyang, and Kerstin Wolk. "Therapeutic opportunities of the IL-22-IL-22R1 system." *Nature reviews Drug discovery* 13.1 (2014): 21-38.
- Schaefer, Laura, et al. "The antimicrobial compound reuterin (3-hydroxypropionaldehyde) induces oxidative stress via interaction with thiol groups." *Microbiology* 156.6 (2010): 1589-1599.
- Sheehan, Michelle M., et al. "Identification and characterization of a lysis module present in a large proportion of bacteriophages infecting *Streptococcus thermophilus*." *Applied and environmental microbiology* 65.2 (1999): 569-577.
- Shi, Yibo, et al. "Characterization and determination of holin protein of *Streptococcus suis* bacteriophage SMP in heterologous host." *Virology journal* 9.1 (2012): 70.
- Singh, Kailash, et al. "Interleukin-35 administration counteracts established murine type 1 diabetes—possible involvement of regulatory T cells." *Scientific reports* 5 (2015).
- Sovran, Bruno, et al. "IL-22-STAT3 pathway plays a key role in the maintenance of ileal homeostasis in mice lacking secreted mucus barrier." *Inflammatory bowel diseases* 21. (2015): 531-542.
- Spinler, Jennifer K., et al. "Human-derived probiotic Lactobacillus reuteri demonstrate antimicrobial activities targeting diverse enteric bacterial pathogens." *Anaerobe* 14.3 (2008): 166-171.
- Steidler, Lothar, et al. "Treatment of murine colitis by Lactococcus lactis secreting interleukin-10." *Science* 289.5483 (2000): 1352-1355.
- Sulakvelidze, Alexander, Zempira Alavidze, and J. Glenn Morris. "Bacteriophage therapy." *Antimicrobial agents and chemotherapy* 45.3 (2001): 649-659.
- Summers WC. Bacteriophage therapy. *Annu Rev Microbiol.* (2001); 55:437-51.
- Sun, Jia, et al. "Pancreatic β -cells limit autoimmune diabetes via an immunoregulatory antimicrobial peptide expressed under the influence of the gut microbiota." *Immunity* 43.2 (2015): 304-317.
- Sznol, Mario, et al. "Use of preferentially replicating bacteria for the treatment of cancer." *Journal of Clinical Investigation* 105.8 (2000): 1027.
- Talarico, T. L., et al. "Production and isolation of reuterin, a growth inhibitor produced by Lactobacillus reuteri." *Antimicrobial agents and chemotherapy* 32.12 (1988): 1854-1858.
- Tannock, Gerald W., et al. "Resource partitioning in relation to cohabitation of Lactobacillus species in the mouse forestomach." *The ISME journal* 6.5 (2012): 927-938.
- Thomas, Carissa M., et al. "Histamine derived from probiotic Lactobacillus reuteri suppresses TNF via modulation of PKA and ERK signaling." *PLoS one* 7.2 (2012): e31951.
- Van Pijkeren, Jan Peter, and Robert A. Britton. "Precision genome engineering in lactic acid bacteria." *Microbial cell factories* 13.1 (2014): S10.
- Van Pijkeren, Jan Peter, et al. "A novel Listeria monocytogenes-based DNA delivery system for cancer gene therapy." *Human gene therapy* 21.4 (2010): 405-416.
- Van Pijkeren, Jan-Peter, and Robert A. Britton. "High efficiency recombinering in lactic acid bacteria." *Nucleic acids research* 40.10 (2012): e76-e76.
- Van Pijkeren, Jan-Peter, et al. "Exploring optimization parameters to increase ssDNA recombinering in Lactococcus lactis and Lactobacillus reuteri." *Bioengineered* 3.4 (2012): 209-217.
- Wang, Ing-Nang, David L. Smith, and Ry Young. "Holins: the protein clocks of bacteriophage infections." *Annual Reviews in Microbiology* 54.1 (2000): 799-825.
- Wang, Xiaoting, et al. "Interleukin-22 alleviates metabolic disorders and restores mucosal immunity in diabetes." *Nature* 514.7521 (2014): 237-241.
- Wells, Jeremy M., et al. "Lactococcus lactis: high-level expression of tetanus toxin fragment C and protection against lethal challenge." *Molecular microbiology* 8.6 (1993): 1155-1162.
- Zhang, Yingnan, et al. "Identification of a small peptide that inhibits PCSK9 protein binding to the low density lipoprotein receptor." *Journal of Biological Chemistry* 289.2 (2014): 942-955.
- Zhou, You, et al. "PHAST: a fast phage search tool." *Nucleic acids research* 39.suppl_2 (2011): W347-W352.

* cited by examiner




FIG. 1A




FIG. 1B




FIG. 2




FIG. 3A




FIG. 3B

In-vivo* assessment of (recombinant) *L. reuteri

Group #1: Sham gavage

#2: LR wild-type (10^9 /day for 7 days)

#3: LR(mIL-22) (10^9 /day for 7 days)




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9A




FIG. 9B




FIG. 9C




FIG. 10




FIG. 11A




FIG. 11B




FIG. 12A




FIG. 12B

1

**METHODS FOR SYSTEMICALLY
DELIVERING POLYPEPTIDES AND
MICROORGANISMS THEREFOR**

STATEMENT REGARDING FEDERALLY
SPONSORED RESEARCH

This invention was made with government support under CA101573 and CA102948 awarded by the National Institutes of Health. The government has certain rights in the invention.

FIELD OF THE INVENTION

The invention is directed to systemically delivering polypeptides to the bloodstream of a subject, such as through administering into the gastrointestinal tract of the subject a microorganism that produces and releases the polypeptides. The invention is also directed to microorganisms suitable for this purpose.

BACKGROUND

Polypeptides, such as enzymes, antibodies, hormones, cytokines, etc., are tremendously useful as therapeutic agents. However, routes for systemically introducing such polypeptides to a subject are limited. Oral administration of the polypeptides is typically not feasible, as the polypeptides are either degraded in the gastrointestinal tract or are blocked from reaching the bloodstream. Direct intravenous administration is therefore the major route by which polypeptides are systemically introduced.

Certain types of genetically engineered bacteria have been used as vehicles for locally delivering polypeptides to various tissues. Engineered *Lactococcus lactis*, for example, has been administered intragastrically for delivering polypeptides such as trefoil factors and interleukin-10 locally to intestinal/mucosal tissues. See Steidler et al. 2000 and Huyghebaert et al. 2005. However, a systemic increase in polypeptides delivered via *Lactococcus lactis* was not found.

Other types of genetically engineered bacteria have been used as vehicles for delivery of polypeptides to tumors in the body. An engineered *Bifidobacterium* strain, for example, has been shown to translocate from the gastrointestinal tract after oral administration and target to, replicate in, and express genes within tumors. See Cronin et al. 2010. This effect, however, depends on the unique ability of the *Bifidobacterium* to translocate from the gastrointestinal tract to extra-intestinal sites in the body. While *Bifidobacterium* may serve as a useful delivery vehicle for some purposes, the systemic distribution of the *Bifidobacterium* is potentially deleterious in certain subject populations such as immunocompromised patients.

Engineered bacteria capable of being administering into the gastrointestinal tract and delivering polypeptides in the bloodstream without systemic levels of the bacteria themselves being increased are needed.

SUMMARY OF THE INVENTION

The invention is directed to methods and microorganisms for systemically introducing a polypeptide in a bloodstream of a subject.

One method comprises administering into the gastrointestinal tract of the subject a bacterium configured to produce and release the polypeptide. The bacterium may com-

2

prise a recombinant gene configured to express the polypeptide. The bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream of the subject, preferably in a detectable amount.

5 In some versions, the bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream of the subject without the bacterium being substantially introduced in the bloodstream of the subject.

10 In some versions, the bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream in an amount effective to induce at least one direct systemic effect in the subject. In some versions, the bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream in an amount effective to induce at least one direct effect in a non-gastrointestinal tissue in the subject. In some versions, the bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream in an amount effective to induce at least one direct effect in a tissue selected from the group consisting of liver, muscles, lungs, kidneys, pancreas, and adipose tissue in the subject.

15 In some versions, the subject suffers from a condition treatable with systemic introduction of the polypeptide. In some versions, the subject suffers from a condition treatable with systemic introduction of the polypeptide but not treatable with local introduction of the polypeptide to the gastrointestinal tract without systemic introduction of the polypeptide. In either case, the polypeptide is introduced in the bloodstream of the subject in an amount effective to treat the condition, independent of the bacterium getting into the bloodstream.

20 In some versions, the polypeptide is a therapeutic polypeptide.

25 In some versions, the polypeptide is selected from the group consisting of a cytokine, a hormone, an antibody, an antimicrobial peptide, and an antigenic peptide.

30 In some versions, the polypeptide is selected from the group consisting of IL-22, IL-35, insulin, leptin, cathelicidin related antimicrobial peptide, a peptide inhibitor of PCSK9, and an endolysin.

35 In some versions, the subject suffers from at least one condition selected from the group consisting of insulin resistance, hyperglycemia, lipid dysregulation, hyperlipidemia, and obesity, and wherein the polypeptide is introduced in the bloodstream of the subject in an amount effective to treat the at least one condition.

40 In some versions, the bacterium comprises a bacterium other than a member of the *Bifidobacterium* genus. In some versions, the bacterium comprises a member of lactic acid bacteria, such as a member of lactic acid bacteria other than a member of the *Lactococcus* genus. In some versions, the bacterium comprises a member of *Lactobacillus*, such as *Lactobacillus reuteri*.

45 A microorganism of the invention comprises a bacterium comprising a recombinant gene configured to express a polypeptide, wherein the bacterium is configured to produce and release the polypeptide and is capable of introducing the polypeptide in the bloodstream of the subject.

50 In some versions, the bacterium is capable of introducing the polypeptide in the bloodstream of the subject without the bacterium being substantially introduced in the bloodstream of the subject.

55 In some versions, the polypeptide is capable of treating a condition in a subject with systemic introduction of the polypeptide in the subject.

In some versions, the polypeptide is selected from the group consisting of IL-22, IL-35, insulin, leptin, cathelicidin related antimicrobial peptide, a peptide inhibitor of PCSK9, and an endolysin.

In some versions, the bacterium comprises a bacterium other than a member of the *Bifidobacterium* genus. In some versions, the bacterium comprises a member of lactic acid bacteria, such as a member of lactic acid bacteria other than a member of the *Lactococcus* genus. In some versions, the bacterium comprises a member of *Lactobacillus*, such as *Lactobacillus reuteri*.

The objects and advantages of the invention will appear more fully from the following detailed description of the preferred embodiment of the invention made in conjunction with the accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A and 1B show mutation rates of various types of bacteria.

FIG. 2 is a plasmid map of the pVPL3461 murine interleukin-22 (mIL-22)-expressing plasmid of the invention, showing an erythromycin resistance gene (Em), an chloramphenicol resistance gene coding sequence (Cm), a phelp promoter (Phelp) (Riedel et al. 2007) for the chloramphenicol resistance gene coding sequence, an mIL-22 coding sequence (mIL-22), a signal peptide for secretion of mIL-22 (SP), a promoter for the signal peptide and the mIL-22 coding sequence (Promoter), and an inverted repeat (IR), which serves as a transcriptional terminator.

FIGS. 3A and 3B show secretion of mIL-22 from engineered *L. reuteri* cells. FIG. 3A shows mIL-22 secretion from *L. reuteri* cells harboring the pVPL3461 plasmid compared to wild-type *L. reuteri* cells as a control. FIG. 3B shows mIL-22 secretion from *L. reuteri* cells harboring a chromosomal copy of a mIL-22 gene compared to wild-type *L. reuteri* cells as a control.

FIG. 4 shows a schema of methods for assessing delivery of mIL-22 to mice from orally administered *L. reuteri* cells harboring the pVPL3461 plasmid, including detecting plasma mIL-22 levels and determining expression of mIL-22 target genes reg3-beta and reg3-gamma.

FIG. 5 shows plasma IL-22 levels in sham gavaged mice (untreated), mice gavaged with wild-type *L. reuteri* (1×10^9 CFU), and mice gavaged with *L. reuteri* engineered to secrete IL-22 (1×10^9 CFU). Eight-week-old male C57BL/6 mice (n=8/group) were gavaged daily for 7 days. Blood was collected from the animals prior to gavage treatment (T=0) and one hour after gavage at the 7th day of administration (T=7). Center lines show the median values. Box limits indicate the 25th and 75th percentiles as determined by R software. Whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles.

FIG. 6 shows counts of bacteria detected in blood from the same mice described above for FIG. 5.

FIG. 7 shows jejunal expression of mIL-22 target genes reg3-beta (reg3B) and reg3-gamma (reg3G) in the same mice described above for FIG. 5. After 7 days gavage, animals were sacrificed, and part of the small intestine (jejunum) was subjected to total RNA isolation followed by cDNA synthesis and real-time PCR. Fold changes and significance are reported based on comparison to the untreated group, and data is normalized against the house-keeping gene 1-actin. Data were analyzed with the REST software package (Pfaffl et al. 2002). Data are presented in a box-whisker plot (see comments above with respect to FIG. 5 for details).

FIG. 8 shows liver expression of lipopolysaccharide-binding protein (LBP) in the same mice described above for FIG. 5.

FIGS. 9A-9C show results from length measurements in animals after eight weeks of high-fat diet feeding (T0) and after seven subsequent weeks of treatment (T7) of daily sham gavage of PBS without bacteria (sham), daily gavage of *L. reuteri* VPL1014 (LR), or daily gavage of the IL-22-secreting *L. reuteri* VPL3461 (LR-IL22). FIG. 9A shows length measurements at T0. FIG. 9B shows growth at T7. FIG. 9C shows length measurements of live versus dead animals.

FIG. 10 shows growth hormone levels in the serum of the mice described above for FIGS. 9A-9C at T7.

FIGS. 11A and 11B show percentage differences in body mass index (BMI) in the mice described above for FIGS. 9A-9C. FIG. 11A shows differences in BMI over the course of seven weeks of treatment (T7-T0). FIG. 11B shows differences in BMI over the course of six weeks of treatment (T7-T1, wherein T1 refers to the time after one week of treatment).

FIGS. 12A and 12B show absolute liver weights and liver weights relative to mouse body weights in the mice described above for FIGS. 9A-9C at T7.

DETAILED DESCRIPTION OF THE INVENTION

The invention provides microorganisms such as bacteria that are capable of introducing polypeptides in the bloodstream of a subject. The invention provides microorganisms such as bacteria that are more specifically capable of introducing the polypeptide in the bloodstream of the subject without the microorganism itself being substantially introduced in the bloodstream of the subject. The invention also provides microorganisms such as bacteria that are capable of introducing the polypeptide systemically in the subject without the microorganism itself being substantially introduced systemically in the subject. "Introduce" and its grammatical equivalents refer to delivery to a site in the body. The introducing may result in detectable presence at that site. "Systemically introduce" and its grammatical equivalents refer to delivery to the bloodstream or sites in the body via the bloodstream. The systemic introducing may result in detectable presence in the bloodstream or such sites. The sites in which the polypeptides are systemically introduced include sites or tissues perfused with the bloodstream and which are permeable to polypeptides. The sites in which the polypeptides are systemically introduced include sites or tissues other than those in the gastrointestinal tract. Exemplary sites or tissues include the liver, muscles, lungs, kidneys, pancreas, adipose tissue, or any other site or tissue in the body.

The bacteria of the invention include certain commensal or probiotic bacteria. The bacteria may include non-pathogenic, Gram-positive bacteria capable of anaerobic growth. The bacteria are preferably viable in the gastrointestinal tract of mammals. The bacteria may be food grade.

Exemplary bacteria include species of lactic acid bacteria (i.e., species of the order Lactobacillales). The bacteria may include species of lactic acid bacteria other than species of the *Lactococcus* genus. The bacteria may include species other than species of the *Bifidobacterium* genus. Exemplary bacteria more preferably include species of the *Lactobacillus* genus.

Exemplary species from the *Lactobacillus* genus include *L. acetototerans*, *L. acidifarinae*, *L. acidiphiscis*, *L. acidophi-*

lus, *L. agilis*, *L. algidus*, *L. atimentarius*, *L. amytoliticus*, *L. amylophilus*, *L. amylophilus*, *L. amylophorus*, *L. animatis*, *L. antri*, *L. apodemi*, *L. aviarius*, *L. bifermentans*, *L. brevis*, *L. buchneri*, *L. camelliae*, *L. casei*, *L. cateniformis*, *L. ceti*, *L. coleohominis*, *L. collinoides*, *L. composti*, *L. concavus*, *L. coryniformis*, *L. crispatus*, *L. crustorum*, *L. curvatus*, *L. delbrueckii* subsp. *delbrueckii*, *L. delbrueckii* subsp. *butgaricus*, *L. delbrueckii* subsp. *lactis*, *L. dextrinicus*, *L. dioliv-orans*, *L. equi*, *L. equigenerosi*, *L. farraginis*, *L. farciminis*, *L. fermentum*, *L. fornicalis*, *L. fructivorans*, *L. frumenti*, *L. fuchuensis*, *L. gallinarum*, *L. gasseri*, *L. gastricus*, *L. ghanensis*, *L. graminis*, *L. hammesii*, *L. hamsteri*, *L. harbinensis*, *L. hayakitensis*, *L. helveticus*, *L. hitgardii*, *L. homohiochii*, *L. iners*, *L. ingluviei*, *L. intestinalis*, *L. jensenii*, *L. johnsonii*, *L. katixensis*, *L. kefiranofaciens*, *L. kefirii*, *L. kimchii*, *L. kitasatonis*, *L. kunkeei*, *L. leichmannii*, *L. lindneri*, *L. malefermentans*, *L. mati*, *L. manihotivorans*, *L. mindensis*, *L. mucosae*, *L. murinus*, *L. nagelii*, *L. namurensis*, *L. nantensis*, *L. oligofermentans*, *L. oris*, *L. panis*, *L. pantheris*, *L. parabrevis*, *L. parabuchneri*, *L. paracollinoides*, *L. parafarraginis*, *L. parakefirii*, *L. parati-mentarius*, *L. paraplantarum*, *L. pentosus*, *L. perolens*, *L. plantarum*, *L. pontis*, *L. psittaci*, *L. renmini*, *L. reuteri*, *L. rhamnosus*, *L. rimae*, *L. rogosae*, *L. rossiae*, *L. ruminis*, *L. saerimneri*, *L. sakei*, *L. salivarius*, *L. sanfranciscensis*, *L. satsumensis*, *L. secaliphilus*, *L. sharpeae*, *L. siliginis*, *L. spicheri*, *L. suebicus*, *L. thailandensis*, *L. ultunensis*, *L. vaccिनostericus*, *L. vaginalis*, *L. versmoldensis*, *L. vini*, *L. vitulinus*, *L. zaeae*, and *L. zymae*.

A particularly preferred bacterium is *L. reuteri*.

A bacterium that can bind mucus in the gastrointestinal tract is preferred but not required. We surmise that binding mucus in the gastrointestinal tract may place the bacterium in close proximity to epithelial cells. Release of polypeptides so close to the epithelial cells may help systemic delivery of the polypeptides. A bacterium capable of binding mucus is *L. reuteri*, such as *L. reuteri* VPL1014, discussed below in the examples. The ability to bind mucus can be mediated by the presence of a mucus-binding protein, such as the cell-mucus binding protein CmbA (Jensen et al. 2014).

The bacterium preferably has a low mutation rate. The bacterium preferably has a mutation rate of less than about 100×10^{-10} mutations per cell per generation, less than about 60×10^{-10} mutations per cell per generation, and more preferably less than about 20×10^{-10} mutations per cell per generation.

The bacterium may be engineered to express a polypeptide of interest. The bacterium accordingly may comprise a recombinant gene configured to express the polypeptide of interest. The bacterium may alternatively or additionally comprise a recombinant DNA sequence that results in increased expression of the polypeptide of interest. “Recombinant” used in reference to a gene refers herein to a sequence of nucleic acids that are not naturally occurring in the genome of the bacterium. The non-naturally occurring sequence may include a recombination, substitution, deletion, or addition of one or more bases with respect to the nucleic acid sequence originally present in the natural genome of the bacterium. “Gene” refers to the collection of genetic elements involved in expressing a coding sequence and may include, in addition to the coding sequence, a promoter, a ribosomal binding site, an enhancer, etc. In some versions, increased expression of the polypeptide of interest can result from introducing or modifying (e.g., recombining, substituting, deleting, etc.) genes or other genetic elements

responsible for regulating expression of the polypeptide of interest, such as genes for transcription factors or signaling factors.

The bacterium may be engineered to produce and release the polypeptide. As used herein, “release” used with respect to the bacterium releasing the polypeptide refers to disposing the polypeptide outside the bacterium, i.e., in the extracellular environment of the microorganism. Release may occur through secretion of the polypeptide or through lysis of the bacterium, among other possible mechanisms. Elements for engineering a bacterium to secrete a polypeptide are well known in the art. Typical elements include a signal peptide-encoding sequence placed upstream of—and in-frame with—the coding sequence of the polypeptide to be secreted. The sequences of a large number of signal peptides for bacteria are known in the art. Exemplary signal peptide sequences are available at www.cbs.dtu.dk/services/SignalP/. Elements for inducing a bacterium to lyse include lytic proteins, which can be expressed from a bacterium through recombinant engineering. As used herein, “lytic protein” refers to any protein that causes or aids in the lysis of a microorganism.

Lytic proteins are well known in the art. A number of lytic proteins, for example, are found in bacteriophages and serve to lyse microorganisms during the lytic stages of the bacteriophage’s life cycle. These include holins and lysins (Sheehan et al. 1999). During bacteriophage replication, biologically active lysins are present in the cytosol but require expression of a membrane protein, holin, to release the virions from the cell. When holin levels are optimal, the lysin can access the peptidoglycan layer for cleavage which leads to bacterial cell lysis (Wang et al. 2000). So far, five main groups of lysins have been identified that can be distinguished from one and another based on the cleavage specificity of the different bonds within the peptidoglycan (Fischetti 2009). Structurally, lysins can consist of a single catalytic domain, which generally is typical for lysins derived from bacteriophages targeting Gram-negative bacteria (Cheng et al. 1994). Bacteriophages targeting Gram-positive bacteria typically encode lysins that contain multiple domains: a N-terminal catalytic domain and a C-terminal cell-wall binding domain (Nelson et al. 2006, Navarre et al. 1999). A few lysins have been identified that have three domains (Becker et al. 2009).

A number of other lytic proteins are native to the microorganisms themselves (Feliza et al. 2012, Jacobs et al. 1994, Jacobs et al. 1995, López et al. 1997). These lytic proteins may affect cell wall metabolism or introduce nicks in the cell wall. Five protein classes are differentiated by the wall component they attack (Loessner et al. 2005, Loessner et al. 2002).

In some versions of the invention, the microorganism is configured to constitutively express a lysin and to express a holin in a maltose-dependent manner. In some versions, the microorganism is configured to express both a lysin and a holin in a maltose-dependent manner.

Lytic proteins can be expressed in a maltose-dependent manner by operably connecting the coding sequence of the lytic protein to a maltose-sensitive promoter. “Coding sequence” refers to a nucleic acid in a gene that encodes the gene product. “Promoter” refers to any nucleic acid that confers, activates, or enhances expression of an operably connected coding sequence. “Operably connected” generally refers to a connection of two genetic elements in a manner wherein one can operate on or have effects on the other. “Operably connected” used in reference to a promoter and a coding sequence refers to a connection between the

promoter and the coding sequence such that the coding sequence is under transcriptional control of the promoter. For example, promoters are generally positioned 5' (upstream) of a coding sequence to be operably connected to the promoter. In the construction of heterologous promoter/ coding sequence combinations, it is generally preferred to position the promoter at a distance from the transcription start site that is approximately the same as the distance between that promoter and the coding sequence it controls in its natural setting, i.e. in the gene from which the promoter is derived. As is known in the art, some variation in this distance can be accommodated without loss of promoter function.

Operably connecting a maltose-inducible promoter to the coding sequence of a lytic protein induces lysis of the microorganism, release of the lytic protein, and release of any other polypeptides made by the microorganism, in a maltose-dependent manner. Such release occurs in the gastrointestinal tract due to natural levels of maltose therein.

An exemplary maltose-inducible promoter is represented by SEQ ID NO: 1, which is a maltose-inducible promoter found in *L. reuteri*. The maltose-inducible promoter represented by SEQ ID NO: 1 or variants thereof are suitable for use in the present invention. Variants of SEQ ID NO:1 include sequences at least about 80% identical, at least about 83% identical, at least about 85% identical, at least about 87% identical, at least about 90% identical, at least about 83% identical, at least about 95% identical, at least about 97% identical, at least about 98% identical, or at least about 99% identical to SEQ ID NO: 1 Other methods of inducing lysis of bacteria in vivo are known.

The bacteria can be engineered using any methods known in the art. General methods are provided in Green et al. 2012. Methods for engineering lactic acid bacteria such as *L. lactis* are provided by van Pijkeren et al. 2012, Oh et al. 2014, and Barrangou et al. 2016.

The recombinant gene may be incorporated into the chromosome of the bacterium or may be included on an extra-chromosomal plasmid. The extra-chromosomal plasmid may replicate at any copy number in the cell and, accordingly, be a single-copy plasmid, a low-copy plasmid, or a high-copy plasmid. The extra-chromosomal plasmid is preferably substantially stable within the bacterium. The rate of loss of the extra-chromosomal plasmid from the bacterium is preferably less than about 10% per generation, less than about 5% per generation, or less than about 1% per generation, wherein percent per generation refers to the percent of the population per generation in which the plasmid is lost.

The bacterium may be engineered to produce and release any polypeptide of interest. The polypeptide may have any of a number of amino acid chain lengths. In some versions, the polypeptide may have an amino acid chain length of from about 2 to about 4000 amino acids, from about 2 to about 3000 amino acids, from about 2 to about 2000 amino acids, from about 2 to about 1500 amino acids, from about 2 to about 1000 amino acids, from about 2 to about 500 amino acids, from about 3 to about 250 amino acids, or from about 3 to about 225 amino acids. The polypeptide may have a net positive charge at neutral pH, a net negative charge at neutral pH, or a net neutral charge at neutral pH. The polypeptide is preferably soluble in water. The polypeptide may form a globular or fibrous structure or may have an intrinsically disordered structure.

The polypeptide may have any of a number of functionalities. The polypeptide, for example, may be enzymatic or non-enzymatic. The polypeptide may be fluorescent or non-

fluorescent. Within the physiological context of a mammal, the polypeptide may be a cytokine, a hormone, an antibody, an antimicrobial peptide, and an antigenic peptide, among others.

Exemplary classes of cytokines include interleukins, lymphokines, monokines, interferons (IFNs), colony stimulating factors (CSFs), among others. Specific exemplary cytokines include IL-1 alpha (IL1a), IL-1 beta (IL1b), IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, IL-21, IL-22, IL-23, IL-24, IL-25, IL-26, IL-27, IL-28, IL-29, IL-30, IL-31, IL-32, IL-33, IL-35, IL-36, IFN-alpha, IFN-beta, IFN-gamma, TNF-alpha, TNF-beta, CNTF (C-NTF), LIF, OSM (oncostatin-M), EPO (erythropoietin), G-CSF (GCSF), GM-CSF (GMCSF), M-CSF (MCSF), SCF, GH (growth hormone), PRL (prolactin), aFGF (FGF-acidic), bFGF (FGF-basic), INT-2, KGF (FGF7), EGF, TGF-alpha, TGF-beta, PDGF, betacellulin (BTC), SCDGF, amphiregulin, and HB-EG, among others.

Exemplary hormones include epinephrine, melatonin, triiodothyronine, thyroxine, amylin (or islet amyloid polypeptide), adiponectin, adrenocorticotrophic hormone (or corticotropin), angiotensinogen, angiotensin, antidiuretic hormone (or vasopressin, arginine vasopressin), atrial-natriuretic peptide (or atriopeptin), brain natriuretic peptide, calcitonin, cholecystokinin, corticotropin-releasing hormone, cortistatin, enkephalin, endothelin, erythropoietin, follicle-stimulating hormone, galanin, gastric inhibitory polypeptide, gastrin, ghrelin, glucagon, glucagon-like peptide-1, gonadotropin-releasing hormone, growth hormone-releasing hormone, hepcidin, human chorionic gonadotropin, human placental lactogen, growth hormone, inhibin, insulin, insulin-like growth factor (or somatomedin), leptin, lipotropin, luteinizing hormone, melanocyte stimulating hormone, motilin, orexin, oxytocin, pancreatic polypeptide, parathyroid hormone, pituitary adenylate cyclase-activating peptide, prolactin, prolactin releasing hormone, relaxin, renin, secretin, somatostatin, thrombopoietin, thyroid-stimulating hormone (or thyrotropin), thyrotropin-releasing hormone, and vasoactive intestinal peptide, among others.

Other physiologically active peptides include glucagon-like peptide-1 (GLP-1); tachykinin peptides, such as substance P, kassinin, neurokinin A, eledoisin, and neurokinin B; peptide PHI 27 (peptide histidine isoleucine 27); pancreatic polypeptide-related peptides, such as NPY (neuropeptide Y), PYY (peptide YY), and APP (avian pancreatic polypeptide); opioid peptides, such as proopiomelanocortin (POMC) peptides and prodynorphin peptides; AGG01; B-type natriuretic peptide (BNP); lactotriptides; and peptides that inhibit PCSK9 (Zhang et al. 2014).

Exemplary antibodies include single-chain antibodies, single-domain antibodies (sdAbs), and single-chain variable fragments (scFvs).

Exemplary antimicrobial peptides include cathelicidins, defensins, protegrins, mastoparan, poneratoxin, cecropin, moricin, melittin, magainin, dermaseptin, and others.

In preferred versions, the polypeptide that is systemically introduced is a polypeptide capable of treating a condition in a subject with its systemic introduction. In some versions, the polypeptide that is systemically introduced is a polypeptide capable of treating a condition in a subject with its systemic introduction but is not capable of treating a condition in the subject with its local introduction to the gastrointestinal tract alone. The condition may be any condition described herein.

The inventors have unexpectedly found that administering *L. reuteri* to the gastrointestinal tract is capable of

delivering produced polypeptides to the bloodstream without the bacteria themselves being introduced in the bloodstream. Accordingly, an aspect of the invention includes administering an amount of a bacterium of the invention into the gastrointestinal tract of a subject. The bacterium may be administered in any amount effective to introduce the polypeptide in the bloodstream of the subject. Exemplary amounts include from about 1×10^3 to about 1×10^{15} , from about 1×10^5 to about 1×10^{13} , from about 1×10^7 to about 1×10^{11} , or about 1×10^9 colony forming units (CFU). Amounts above and below these ranges may be acceptable.

The bacterium can be administered to the gastrointestinal tract by any method known in the art. The bacterium may be administered orally, rectally, or directly into the gastrointestinal tract via a stoma. The bacterium is preferably administered directly into or upstream of the small intestines, so that the bacterium ultimately passes through or into the small intestines. The bacterium may be swallowed or introduced via a tube.

The bacterium may be combined in a composition with a pharmaceutically acceptable excipient, carrier, buffer, stabilizer or other material well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the bacterium. The precise nature of the carrier or other material may depend on the route of administration. The composition may be liquid, solid, or semi-solid. The composition may comprise a foodstuff or may take the form of a pharmaceutical composition. Those of relevant skill in the art are well able to prepare suitable compositions.

The subject to which the bacterium is administered may be an animal, such as a mammal or, more specifically, a human.

In some versions of the invention, the bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream in an amount effective to induce at least one systemic effect in the subject, such as at least one effect in a non-gastrointestinal tissue in the subject. As used herein, "systemic effect" refers to an effect that occurs at a site or tissue in the body other than where the polypeptide is initially released from the bacterium. In the present case, the term refers to an effect that occurs at a site or tissue other than the gastro-intestinal tract, such as the liver, muscles, lungs, kidneys, pancreas, adipose tissue, or others. The systemic effect preferably occurs by virtue of the polypeptide being systemically introduced and accessing a site or tissue other than gastrointestinal tissue via bloodstream, such that the effect is a direct effect of the polypeptide and is not a secondary effect of a primary effect of the polypeptide in the gastrointestinal tissue. Such effects are referred to herein as "direct systemic effects." Direct systemic effects of the systemically introduced polypeptide can be distinguished from secondary effects, for example, by comparing effects resulting from administering the polypeptide-producing bacterium into the gastrointestinal tract with effects resulting from systemically administering the polypeptide directly into the bloodstream and effects resulting from locally administering the polypeptide directly into the gastrointestinal tract. Direct systemic effects are those that are mirrored by the systemic administration of the polypeptide into the bloodstream but not the local administration of the polypeptide into the gastrointestinal tract. The presence of direct systemic effects of the polypeptide resulting from administering the polypeptide-producing bacterium into the gastrointestinal tract can be an indicator of the polypeptide entering the bloodstream, whether or not the polypeptide itself is detected in the bloodstream.

In some versions, the bacterium is administered to a subject that suffers from a condition treatable with systemic introduction of the polypeptide. In yet other versions, the bacterium is administered to a subject that suffers from a condition treatable with systemic introduction of the polypeptide but is not treatable with only local introduction of the polypeptide to the gastrointestinal tract. In either case, the polypeptide is introduced in the bloodstream of the subject in an amount effective to treat the condition. As used herein, "treat" used in reference to a condition refers to ameliorating to any extent the condition itself or any symptom associated therewith.

In some versions, the bacterium is administered to a subject to prevent or inhibit the development of any of the conditions or associated symptoms described herein. The subject in such a case may show early signs of the condition or symptom, have a genotype that predisposes the subject to develop the condition or symptom, have a behavior or environmental situation that predisposes the subject to develop the condition or symptom, or otherwise be predisposed to develop the condition or symptom.

A particular aspect of the invention is directed to introducing interleukin-22 (IL-22) in the bloodstream of a subject by administering a bacterium comprising a recombinant interleukin-22 (IL-22) gene. The IL-22 may be introduced in the bloodstream of the subject in an amount effective to induce at least one IL-22-dependent effect in a non-gastrointestinal tissue in the subject. An exemplary IL-22 that can be systemically introduced comprises a sequence of SEQ ID NO:2 or a sequence at least about 90%, 95%, 97%, 99% or more identical thereto.

IL-22 is capable of inducing a number of effects in non-gastrointestinal tissues when circulating systemically through the bloodstream. In respiratory epithelial cells, for example, IL-22 can increase antibacterial defense, elevate mucus production, enhance proliferation, and raise production of granulocyte-attracting chemokines. In synovial fibroblasts, IL-22 can elevate RANKL expression and increase production of monocyte-attracting chemokines. In pancreatic cells, IL-22 can increase protection against damage, inhibit autophagy, and enhance islet proliferation. In hepatocytes, IL-22 can increase acute-phase protein production, increase protection against damage, and elevate liver progenitor cell proliferation. In epidermal keratinocytes, IL-22 can increase antibacterial defense, retard differentiation and cornification, induce production of granulocyte-attracting chemokines, elevate migration and tissue remodeling, and enhance STAT3 and IL-20 expression. See Sabat et al. 2014 and Wang et al. 2014 for additional direct IL-22-dependent effects.

Another particular aspect of the invention is directed to introducing interleukin-22 (IL-22) in the bloodstream of a diabetic subject by administering a bacterium comprising a recombinant interleukin-22 (IL-22) gene. The diabetic subject may suffer from type 1 or type 2 diabetes. IL-22 has been shown to have a number of ameliorative effects in diabetic subjects. In type 2 diabetic subjects, these effects include ameliorating hyperglycemia, insulin resistance, hyperlipidemia, lipid dysregulation in the liver and adipose tissues, endotoxemia, and chronic inflammation. See Wang et al. 2014. As shown in the present examples, IL-22 is also capable of reducing body mass index (BMI).

More generally, subjects in which IL-22 is introduced may include those suffering from of insulin resistance, hyperglycemia, lipid dysregulation, hyperlipidemia, obesity, or other manifestations of metabolic syndrome. The systemic effect of the systemic introduction of IL-22 to such

subjects may include a reduction in body mass index (BMI), liver weight, liver triglycerides, glucose intolerance, and insulin resistance, or other effects described elsewhere herein.

Another polypeptide that may be introduced in the bloodstream of a subject with the bacteria of the invention is interleukin-35 (IL-35). Systemic administration of IL-35 treats type-1 diabetes and inhibits or slows the development of type-1 diabetes. See Singh et al. 2015. An exemplary IL-35 that can be systemically introduced is a human recombinant IL-35 comprising a sequence of SEQ ID NO:3 or a sequence at least about 90%, 95%, 97%, 99% or more identical thereto. Bacteria of the invention comprising a recombinant IL-35 gene can be administered to subjects with diabetes, such as type 1 diabetes, to systemically introduce the IL-35 polypeptide in the bloodstream of the subject in an amount effective to treat the diabetes in the subject.

Another polypeptide that may be introduced in the bloodstream of a subject with the bacteria of the invention is insulin. The insulin can be produced in single-chain form. See, e.g., Rajpal et al. 2009. An exemplary insulin that can be systemically introduced includes the insulin A-chain connected to the insulin B-chain by the linker sequence QRGGGGGQR (SEQ ID NO:4). See Rajpal et al. 2009. Single-chain insulin retains all of the physiological effects of traditional two-chain insulin, including stimulation of glucose uptake into adipocytes, and suppression of hepatic gluconeogenesis. Bacteria of the invention comprising a recombinant insulin gene can be administered to subjects with diabetes, insulin resistance, or hyperglycemia to systemically introduce the insulin polypeptide in the bloodstream of the subject in an amount effective to treat the diabetes, or hyperglycemia in the subject.

Another polypeptide that may be introduced in the bloodstream of a subject with the bacteria of the invention is leptin. Leptin is made by adipose tissue and regulates energy balance by acting on receptors in the brain. Congenital leptin deficiency (CLD), or generalized lipodystrophy results in a lack of leptin and can lead to a litany of disorders, including morbid obesity (in the case of CLD), diabetes, and infertility. Systemic leptin replacement therapy mitigates these disorders. An exemplary leptin polypeptide that can be systemically introduced includes a polypeptide comprising a sequence of SEQ ID NO:5 or a sequence at least about 90%, 95%, 97%, 99% or more identical thereto. Bacteria of the invention comprising a recombinant leptin gene can be administered to subjects with congenital leptin deficiency or generalized lipodystrophy to systemically introduce the leptin polypeptide in the bloodstream of the subject in an amount effective to treat the obesity, diabetes, infertility or any other aspect of the congenital leptin deficiency or generalized lipodystrophy.

Another polypeptide that may be introduced in the bloodstream of a subject with the bacteria of the invention is cathelicidin related antimicrobial peptide (CRAMP). CRAMP is a small peptide produced by pancreatic islets in response to gut microbiota-derived short-chain fatty acids. Synonyms for CRAMP include CAMP, CAP18, Cnlp, FALL39, and MCL. The islet-derived CRAMP maintains immune homeostasis. CRAMP production is defective in

non-obese diabetic mice, leading to inflammation and activation of diabetogenic T-cells and resulting in type 1 diabetes (Sun et al. 2015). This process can be reversed by direct systemic administration of CRAMP. An exemplary CRAMP polypeptide that can be systemically introduced includes a polypeptide comprising a sequence of SEQ ID NO:6 or a sequence at least about 90%, 95%, 97%, 99% or more identical thereto. Bacteria of the invention comprising a recombinant CRAMP gene can be administered to non-obese diabetic subjects to systemically introduce the CRAMP polypeptide in the bloodstream of the subject in an amount effective to treat the diabetes and/or inflammation in these subjects.

Other polypeptides that may be introduced in the bloodstream of a subject with the bacteria of the invention include peptide inhibitors of PCSK9. PCSK9 (proprotein convertase subtilisin/kexin type 9) is a negative regulator of the hepatic low density lipoprotein receptor. Inhibition of PCSK9 results in LDL cholesterol-lowering effects. A number of peptide inhibitors of PCSK9 are known in the art. See Zhang et al. 2014. Any of these polypeptides or others can be introduced in the bloodstream of a subject with the bacteria of the invention. A particularly preferred peptide inhibitor of PCSK9 is referred to as "Pep2-8," which comprises a sequence of SEQ ID NO:7. Bacteria of the invention comprising a recombinant gene configured to express one or more peptide inhibitors of PCSK9 can be administered to subjects with hypercholesterolemia to systemically introduce the peptide inhibitor of PCSK9 in the bloodstream of the subject in an amount effective to treat the hypercholesterolemia.

Other polypeptides that may be introduced in the bloodstream of a subject with the bacteria of the invention include lysins, such as bacteriophage-derived lysins (endolysins), i.e. enzybiotics. One of the largest concerns in 21st century medicine is the development of microbial antibiotic-resistance. Little progress has been made in the discovery and development of novel antibiotics, and bacteriophage-derived lysins (enzybiotics) constitute promising alternatives to antibiotics. The enzybiotics interfere with peptidoglycan cell wall synthesis, mainly of Gram positive bacteria, but do so in a species specific manner. Exemplary lysins that can be systemically introduced include those described in the references cited herein, all of which are incorporated herein by reference. Bacteria of the invention comprising a recombinant gene configured to express one or more lysins can be administered to subjects with sepsis or infection with pathogens such as *S. aureus* to systemically introduce the lysin in the bloodstream of the subject in an amount effective to treat the sepsis or infection.

The elements and method steps described herein can be used in any combination whether explicitly described or not.

All combinations of method steps as used herein can be performed in any order, unless otherwise specified or clearly implied to the contrary by the context in which the referenced combination is made.

As used herein, the singular forms "a," "an," and "the" include plural referents unless the content clearly dictates otherwise.

Numerical ranges as used herein are intended to include every number and subset of numbers contained within that

range, whether specifically disclosed or not. Further, these numerical ranges should be construed as providing support for a claim directed to any number or subset of numbers in

VPL3461. For animal experiments (see below) 100 μ l of the suspension was administered by oral gavage to the corresponding animals.

TABLE 1

Bacterial strains and plasmids used in the present examples.		
Strain or plasmid	Characteristics	Source
<i>L. reuteri</i> VPL1014	A derivative of <i>L. reuteri</i> ATCC PTA 6475, human breast milk isolate	ATCC PTA 6475, U.S. Pat. 7,344,867 to BioGaia AB (Lerum, SE), and van Pijkeren et al. 2012
<i>L. reuteri</i> VPL3461 pJP028	<i>L. reuteri</i> VPL1014 harboring pVPL3461 Em ^R , derivative of pNZ8048 containing promoter from <i>L. reuteri</i> SD2112, signal peptide from <i>L. reuteri</i> JCM1112, and LPXTG (cell wall anchor domain)	Described herein
pVPL3461	Em ^R , derivative of pJP028 omitting LPXTG domain and harboring mIL-22 gene	Described herein

that range. For example, a disclosure of from 1 to 10 should be construed as supporting a range of from 2 to 8, from 3 to 7, from 5 to 6, from 1 to 9, from 3.6 to 4.6, from 3.5 to 9.9, and so forth.

All patents, patent publications, and peer-reviewed publications (i.e., "references") cited herein are expressly incorporated by reference to the same extent as if each individual reference were specifically and individually indicated as being incorporated by reference. In case of conflict between the present disclosure and the incorporated references, the present disclosure controls.

It is understood that the invention is not confined to the particular construction and arrangement of parts herein illustrated and described, but embraces such modified forms thereof as come within the scope of the claims.

EXAMPLES

Bacterial strains, plasmids, media, and culture.

Bacterial strains and plasmids used in the present examples are listed in Table 1. *Lactobacillus reuteri* VPL1014 and its derivatives were routinely cultured at 37° C. in deMan Rogosa Sharpe (MRS) medium (Difco, BD Biosciences). Where appropriate, erythromycin was added to a final concentration of 5 μ g/ml. Competent cells of *L. reuteri* were prepared as described before (Ahrnd et al. 1992). To test IL-22 expression and to prepare bacteria for animal experiments, bacteria were cultured in Lactobacilli Defined Medium-III (LDM-III, Table 2).

Specifically, *L. reuteri* VPL1014 was inoculated in 10 ml MRS broth, and *L. reuteri* VPL3461 was inoculated in 10 ml MRS containing 5 g/ml erythromycin at 37° C. Overnight-cultures of each strain were sub-cultured in MRS at OD₆₀₀=0.1. At OD₆₀₀≥1≤1.2 cells were harvested by centrifugation (5,000 rpm for 5 min), and the cell pellets were washed twice with LDM-III. Washed cell pellets, each derived from 10-ml culture, were stored at -80° C. until use. From this step onwards, no antibiotics were supplemented to the culture media. At the day of gavaging, the cell pellets were resuspended in 10 ml freshly prepared pre-warmed LDM-III and incubated at 37° C. until OD₆₀₀≥3.5≤3.7. Bacteria were concentrated 10-fold, and suspensions were made containing ~10¹⁰ CFU/ml *L. reuteri* VPL1014 or

TABLE 2

Lactobacilli Defined Medium-III (LDM3) composition.		
Basal Medium		
Ingredient	Amount per Liter	
K ₂ HPO ₄	1.50 g	
KH ₂ PO ₄	1.50 g	
Sodium Acetate	15.00 g	
Sodium Citrate, Dihydrate	0.25 g	
Tryptophan	0.05 g	
Asparagine, Monohydrate	0.23 g	
Vitamin-free Casamino acid	10.00 g	
Cysteine-HCl, Monohydrate	0.22 g	
Tween80 (10% v/v)	10 ml	
Dissolve in 937.5 ml of DI water and autoclave at 121° C. for 15 min. Prepare stock solutions (as below) and add to sterile Basal Medium.		
Vitamin Solution (in 25 ml dH ₂ O)		
Thiamin HCl	10 mg	0.5 ml
p-aminobenzoic acid	2 mg	
Calcium pantothenic acid	20 mg	
Niacin	50 mg	
Pyridoxin HCl	25 mg	
Biotin Solution (in 50 ml 0.01M HCl)		
Biotin	4 mg	0.5 ml
96% EtOH	40 μ L	
Riboflavin Solution (in 50 ml dH ₂ O)		
Riboflavin	4 mg	5 ml
Folic Acid Solution (in 50 ml 0.001M NaOH)		
Folic acid	10 mg	0.5 ml
Nucleic Acid Solution (in 15 ml 1M HCl)		
Adenine hemisulfate	50 mg	3 ml
Guanine	40.3 mg	
Cytidylic acid	150 mg	
Uracil Solution (in 10 ml 1M NaOH)		
Uracil	200 mg	1 ml
Thymidine Solution (in 12.5 ml dH ₂ O)		
Thymidine	20 mg	1 ml
Salt Solution (in 10 ml dH ₂ O)		
MgSO ₄	0.793 g	1 ml

15

TABLE 2-continued

Lactobacilli Defined Medium-III (LDM3) composition.		
MnSO ₄	0.128 g	
FeSO ₄	0.130 g	
Glucose (40% w/v)		50 ml
Total 1 L (LDM), final pH6.5		

Mutation Rate.

A suitable microorganism for delivering peptides preferably has a low mutation rate. The mutation rate of *L. reuteri* VPL1014 was compared with that of a number of other microorganisms using conventional methods (Rosche et al. 2000, Foster et al. 2006). As shown in FIGS. 1A and 1B, *L. reuteri* displayed an exceptionally low mutation rate, particularly compared to the other tested probiotic microorganisms.

Reagents and Enzymes.

Cloning was performed via ligation cycle reaction (LCR; Kok et al. 2014). Enzymes and reagents for LCR were purchased from Fermentas. Polymerase chain reaction (PCR) for cloning purposes was performed with the high-fidelity enzyme Phusion Hot Start Polymerase II (Fermentas). PCR for screening purposes was performed with Taq polymerase (Denville Scientific). To concentrate the LCR reaction prior to electrotransformation into *L. reuteri*, we used Pellet Paint Co-Precipitant (Novagen). Oligonucleotides and synthetic double-stranded DNA fragments were purchased from Integrated DNA Technologies. All oligonucleotides and synthetic DNA fragments used in this study are listed in Table 3.

TABLE 3

Oligonucleotides and synthetic DNA used in the present examples.	
Oligo-nucleotides	Sequence
oVPL329	attccttggacttcatttactgggtttaac (SEQ ID NO: 8)
oVPL363	taatgatgagataatgccgactgtac (SEQ ID NO: 9)
oVPL1219	ttcatggggatgaatgcttctgctaatacattaccagttaa tactcgttg (SEQ ID NO: 10)
oVPL1220	cttgggttttctaattttgggttcaaagatcaaacacaagcat tacgtaaaactc (SEQ ID NO: 11)
oVPL1221	gcttgaaacgttcaattgaaatggca (SEQ ID NO: 12)
oVPL1222	tgtaaaaccaataaggactgaagc (SEQ ID NO: 13)
oVPL1223	ggagttgcttcagtccttattgggtttacattcatggggat gaatgcttctgctaataca (SEQ ID NO: 14)
oVPL1224	tgatctttgaacaaaattagaaaaccaaggcttgaacgt tcaattgaaatggcaatta (SEQ ID NO: 15)
oVPL1313	actccctgaagaatataccctcc (SEQ ID NO: 16)

16

TABLE 3-continued

Oligonucleotides and synthetic DNA used in the present examples.	
Oligo-nucleotides	Sequence
oVPL1314	cgctattgagcacagatacagag (SEQ ID NO: 17)
oVPL1315	atgcttccccgtataaccatca (SEQ ID NO: 18)
oVPL1316	ggccatattctgcatcataccag (SEQ ID NO: 19)
oVPL1321	gatcaccgacaagggcctg (SEQ ID NO: 20)
oVPL1322	ggctatgaaactcgtactgcc (SEQ ID NO: 21)
oVPL1325	ggctgtattccccctccatcg (SEQ ID NO: 22)
oVPL1326	ccagttggtaacaatgccatgt (SEQ ID NO: 23)
gVPL1	ATTCATGGGGATGAATGCTTCTGCTAATACATTACCAGTTA ATACTCGTTGTAAATTAGAAGTTAGTAATTTTCAACAACCA TATATTGTTAATCGTACTTTTATGTTAGCTAAAGAAGCTAG TTAGCTGATAATAACTGATGTTTCGTTAATTGGTGAAA AATTATTTTCGTGGTGTAGTGTCTAAAGATCAATGTTATTTA ATGAAACAAGTTTTAAATTTTACTTTAGAAGATGTTTATT ACCACAAAGTGATCGTTTTCAACCATATATGCAAGAAGTTG TTCCATTTTTAACTAAATTAAGTAATCAATTAAGTAGTTGT CATATTAGTGGTGATGATCAAAATATTCAAAAAATGTTTCG TCGTTTAAAAGAACTGTAAAAAATTAGGTGAAAGTGGTG AAATTAAGCTATTGGTGAATTAGATTTATTATTATGAGT TTACGTAATGCTTGTGTTGATCTTTGAACCAAAATTAGAA AACCAAGG (SEQ ID NO: 24)

Construction of *L. reuteri* VPL1014 that Secretes mIL-22.

Our aim was to engineer *Lactobacillus reuteri* VPL1014 to secrete the murine cytokine interleukin-22 (mIL-22). We first opted for expression from the multicopy plasmid pJP028 to maximize mIL-22 production. pJP028 is a derivative of pNZ8048 (de Ruyter et al. 1996) in which the nisin-expression cassette was replaced with a secretion cassette. By PCR (oVPL1221-oVPL1222), we amplified the backbone of pJP028, omitting the cell wall anchor domain, to yield a 4.579 kb product. For optimal expression of mIL-22 in our expression host, *L. reuteri*, we first applied in-silico codon optimization of the mIL-22 coding sequence using the online software, OPTIMIZER (genomes.urv.es/OPTIMIZER/, Table 4) followed by synthesis. The resulting synthetic product (gVPL1) was amplified by PCR (oVPL1219 and oVPL1220), followed LCR (Kok et al. 2014) placing the gVPL1 fragment between the start and stop codon located on the pJP028 backbone. The LCR mixture was precipitated and transformed in *L. reuteri* VPL1014. Transformants were screened by PCR (oVPL329-oVPL363) to confirm cloning of mIL-22. One positive clone was colony purified, a 1.584 kb amplicon was generated by colony PCR, and the integrity of the construct was confirmed by DNA sequencing (GeneWiz). The resultant strain was named VPL3461. We hereafter refer to pVPL3461 when it concerns the plasmid that encodes codon-optimized mIL-22 (FIG. 2).

The nucleotide sequence of pVPL3461 is represented by SEQ ID NO:25. The nucleotide sequence of the IL-22 promoter (*L. reuteri* native promoter) in pVPL3461 is rep-

resented by SEQ ID NO: 26. The nucleotide sequence encoding the signal peptide (SP) in pVPL3461 is represented by SEQ ID NO:27. The nucleotide sequence encoding IL-22 in pVPL3461 is represented by SEQ ID NO:28. The nucleotide sequence of the inverted repeat (IR) in pVPL3461 is represented by SEQ ID NO:29. The nucleotide sequence of the chloramphenicol marker (Cm) in pVPL3461 is represented by SEQ ID NO:30. The nucleotide sequence of the Phelp promoter in pVPL3461 is represented by SEQ ID NO:31. The nucleotide sequence of the erythromycin marker (Em) in pVPL3461 is represented by SEQ ID NO:32.

Additionally, a construct from the pVPL3461 plasmid comprising the promoter, signal peptide, and mIL-22 coding sequence was excised from pVPL3461 and incorporated in the *L. reuteri* chromosome using methods known in the art. The resultant strain was named VPL3461chr.

TABLE 4

Codon optimization table for <i>L. reuteri</i> F275.*#			
Fields: [sequence of nucleotide triplet] [frequency: per thousand] ([number])			
UUU 29.5 (16802)	UCU 9.7 (5521)	UAU 24.8 (14106)	UGU 4.4 (2479)
UUC 11.3 (6447)	UCC 4.0 (2249)	UAC 12.4 (7068)	UGC 1.5 (861)
UUA 36.4 (20706)	UCA 15.0 (8537)	UAA 2.3 (1307)	UGA 0.4 (219)
UUG 15.4 (8765)	UCG 4.3 (2458)	UAG 0.7 (374)	UGG 10.8 (6134)
CUU 22.1 (12555)	CCU 9.7 (5489)	CAU 15.3 (8708)	CGU 14.1 (7995)
CUC 6.3 (3597)	CCC 3.4 (1927)	CAC 8.0 (4562)	CGC 5.9 (3360)
CUA 9.4 (5359)	CCA 17.7 (10090)	CAA 35.0 (19877)	CGA 8.6 (4869)
CUG 5.3 (3005)	CCG 6.0 (3428)	CAG 11.7 (6653)	CGG 9.9 (5610)
AUU 50.7 (28857)	ACU 22.3 (12657)	AAU 36.1 (20541)	AGU 15.6 (8850)
AUC 16.7 (9524)	ACC 9.8 (5550)	AAC 15.8 (8968)	AGC 6.7 (3786)
AUA 5.8 (3300)	ACA 16.6 (9464)	AAA 36.6 (20829)	AGA 3.3 (1872)
AUG 26.9 (15321)	ACG 9.2 (5230)	AAG 30.3 (17232)	AGG 1.4 (792)
GUU 35.0 (19884)	GCU 29.0 (16508)	GAU 43.5 (24740)	GGU 24.3 (13830)
GUC 9.2 (5210)	GCC 12.4 (7053)	GAC 15.2 (8617)	GGC 10.9 (6178)
GUA 16.1 (9176)	GCA 25.3 (14409)	GAA 45.6 (25918)	GGA 19.8 (11244)
GUG 7.7 (4354)	GCG 9.8 (5573)	GAG 11.2 (6360)	GGG 10.1 (5771)

*This table was made based on 568,715 codons among 1,900 CDSs on chromosomal DNA of strain F275.

#Coding GC 39.50% 1st letter GC 51.33% 2nd letter GC 35.17% 3rd letter GC 32.00%.

Determine mIL-22 Secretion.

Strains *L. reuteri* VPL1014 and VPL3461 were cultured in LDM-III as described above, and the supernatants were collected after centrifugation (5 min at 3,214×g), followed by filter-sterilization (0.22 μm, Millipore). One hundred microliters of filter-sterilized supernatant from *L. reuteri* VPL1014 and VPL3461 was assessed for the presence of mIL-22 by ELISA (R&D systems). Production of mIL-22 could not be detected for *L. reuteri* 6575-VPL (15 pg/ml cut-off limit), while strain VPL3461 secreted mIL-22 at levels of 164.2±13.1 ng/ml (n=3). See FIG. 3A.

Secretion of mIL-22 from VPL3461chr was similarly tested. VPL3461chr showed detectable mIL-22 secretion. See FIG. 3B.

Plasmid Stability of pVPL3461 in *L. reuteri* VPL1014.

Prior to assessing biological activity of the *L. reuteri*-produced mIL-22 in mice, we first assessed the stability of pVPL3461. Normally, selection of plasmids is achieved by supplementation of an antibiotic, but we wanted to avoid the supplementation of antibiotics in mice to maintain a fully competent microbiota. VPL3461 was cultured overnight in LDM-III (supplemented with antibiotics). Cells were washed to remove residual antibiotics, followed by sub-culturing to antibiotic-free LDM-III (OD₆₀₀=0.1). After 1 passage (20 hr, ~10 generations), we showed that 96% of the cell population was resistant to erythromycin, demonstrating that the rate of loss of pVPL3461 was 0.4% per generation, and would be considered stable enough for in-vivo assessment of biological activity.

Animal Trial.

Twenty-four 6-week old male B6 mice (C57BL/6J) were purchased from Jackson Labs (Bar Harbor, Me.). Animals were housed at an environment-controlled facility with a 12-hour light and dark cycle. Both diet (standard chow, LabDiet, St Louis, Mo.) and water were freely available to the animals. After transport, animals were allowed to adjust to the new environment for two weeks, after which treatment by gavage started. Three groups of 8 animals per group were treated daily for 7 consecutive days. Treatments were sham gavage where the animals were subjected to insertion of a gavaging needle without administering anything (control), gavage of *L. reuteri* VPL1014 (WT group) and gavage of *L. reuteri* VPL3461 (LR_mIL-22). Bacterial load administered was ~1×10⁹ CFU in a volume of 100 μl of the respective bacterial supernatant. See FIG. 4.

Blood collection to assess plasma IL-22 levels.

At T=0 (prior to the start of treatment) and at T=7 (2 hours after the last gavage) of the animal trial, blood was collected (50 μl per animal) via retro orbital puncture. Plasma was isolated from whole blood sample by centrifugation at 9,000 rpm for 7 min and the plasma fraction was stored at -80° C. until use. By ELISA (as described above) we determined plasma mIL-22 levels. See FIG. 4.

Plasma IL-22 levels after 7 days gavage are shown in FIG. 5. The mice administered *L. reuteri* VPL3461 showed a statistically significant increase in plasma IL-22 levels compared to controls.

We also assessed whether the administered *L. reuteri* VPL1014 and *L. reuteri* VPL3461 could be detected in the bloodstream of the animals after 7 days gavage. From each animal, 50 μl blood was plated on deMan Rogosa Sharp (MRS) medium that is selective for a broad range of lactic acid bacteria, including *L. reuteri*. The results are shown in FIG. 6. The prolonged daily administration of *L. reuteri* did not increase the total number of lactic acid bacteria in the bloodstream. As determined by colony morphology, the bacteria detected in the bloodstream of animals that were gavaged with *L. reuteri* were not *L. reuteri*. *L. reuteri* yields (after 48 h) on MRS plates small-medium sized colonies that are opaque. From the 3 animals in which we detected bacteria in the bloodstream, all colonies were pigmented, mostly yellow, and some were red-ish. Some colonies were also extremely large. The size combined with the pigmented phenotype made it evident that the recovered bacteria in the bloodstream were not *L. reuteri*.

These results indicate that the systemic increase of IL-22 was not a result of *L. reuteri* VPL3461 itself entering the bloodstream.

cDNA Synthesis.

To assess biological functionality of *L. reuteri* secreted mIL-22, we assess gene expression levels of reg3-beta and reg3-gamma. Both genes are known to be upregulated by IL-22 (Loonen et al. 2013, Sovran et al. 2015). Part of the small intestine (jejunum) of each animal was processed for RNA isolation. First, samples were homogenized (Omni TH, Omni International) followed by RNA isolation and on-column DNaseI treatment (Qiagen), after which an additional DNase treatment was conducted (RQI DNase; Promega, Madison, Wis.). RNA was quantified by Qubit analysis (Invitrogen). One μ g RNA was reverse transcribed using the iScript cDNA synthesis kit (Bio-Rad Laboratories, Richmond, Calif.). See FIG. 4.

Quantitative Real-Time PCR.

Relative gene expression levels were determined using the CFX96™ real-time PCR (Bio-Rad). Expression of reg3-beta and reg3-gamma was determined relative to that of the housekeeping gene β -actin. The qRT-PCR was performed with the SYBR Green PCR master mix (Bio-Rad). Primers for amplification of: reg3b (oVPL1313-oVPL1314), reg3g (oVPL1315-oVPL1316), and β -actin (oVPL1325-oVPL1326) are listed in Table 3. Gene expression of the reg genes in the jejunum tissues relative to β -actin was determined by the Relative Expression Software Tool (REST), which allows comparison of gene expression between groups of animals (Pfaffl et al. 2001 and 2002).

As shown in FIG. 7, mice administered IL-22-expressing *L. reuteri* VPL3461 showed an average of 4.7-fold and 3-fold increased expression of reg3-beta and reg3-gamma, respectively in the jejunum compared to mice administered wild-type *L. reuteri*, demonstrating that the secreted IL-22 is biologically active.

We also determined liver expression of the gene encoding the lipopolysaccharide-binding protein (LBP), which is known to be regulated by IL-22. Expression levels were relative to that of the control (β -actin). As shown in FIG. 8, liver LBP expression was not changed in animals that received wild-type *L. reuteri*. Animals administered the IL-22-expressing *L. reuteri* VPL3461 displayed an increased level of LBP gene expression, varying from 1.5-fold to 3.5-fold. We did not detect a statistical difference in gene expression, but we predict including more animals per group will show a statistical difference.

Metabolic Syndrome Trial.

IL-22 has been shown to alleviate metabolic disorders and provide other therapeutic effects in diabetic subjects. See Wang et al. 2014. We tested whether administering IL-22-secreting *L. reuteri* to mice with diet-induced obesity could recapitulate these effects.

Thirty-six 6-week old male B6 mice (C57BL/6J) were purchased from Jackson Labs (Bar Harbor, Me.). Animals were housed at an environmental controlled facility with a 12-hour light and dark cycle. After transport, animals were caged (4 mice per cage) and immediately placed on an ad libitum high-fat diet: 45% kcal fat diet containing 21% milk fat and 2% soybean oil (Cat. No. TD08811, Envigo, Indianapolis, Ind.), for eight weeks. Based on prior work, animals placed on this diet for eight weeks develop signs of metabolic syndrome, including glucose intolerance and insulin sensitivity.

After eight weeks on the high-fat diet, we initiated the treatment of daily gavage for a period of seven weeks. Animals in a first group (12 animals) received a sham

gavage of 100 μ l PBS without bacteria. Animals in a second group (12 animals) received 100 μ l of *L. reuteri* VPL1014 (10^9 CFU). Animals in a third group (12 animals) received 100 μ l of the IL-22-secreting *L. reuteri* VPL3461 (10^9 CFU).

The length (nose to anus) of the animals was determined after eight weeks high-fat diet feeding (T0), and was subsequently determined every week after for seven weeks (T7). Each animal was measured three times and the values were averaged. After eight weeks high-fat diet feeding but prior to the start of the treatment (T0) we observed that the animals assigned to be receiving treatment were similar in length ($P=0.88$) but both groups were marginally smaller than the control group (control vs WT, $P=0.09$; control vs recombinant, $P=0.04$). See FIG. 9A. After seven weeks of treatment (T7) we observed that the animals gavaged with the IL-22-secreting *L. reuteri* grew faster than animals gavaged with *L. reuteri* wild-type ($P<0.0001$) or PBS control ($P<0.0001$). See FIG. 9B. The increased growth is purely driven by recombinant IL-22 that is delivered by *L. reuteri* because *L. reuteri* wild-type does not influence growth compared to the PBS control ($P=0.66$).

Healthy mice have a natural curve in their spine. When mice are obese, the excess weight will press the spine down. When measuring length of the animal from nose to anus, obese mice may be perceived to be longer. To determine if this would have affected our body length data, we measured animals alive and after euthanasia at T7. When anesthetized or dead, any bias derived from differences in curvature will be lost as the animal is completely relaxed. As shown in FIG. 9C, there is no difference in the body length of the animals when measured alive or dead. This finding conclusively confirmed that recombinant *L. reuteri* secreting IL-22 promotes growth.

Our growth data led us to measure growth hormone levels in the serum. As shown in FIG. 10, mice treated with IL-22-secreting *L. reuteri* had increased levels of growth hormone compared to the PBS control ($P=0.03$) at T7. Levels were also higher in the recombinant group compared to the wild-type but this was not statistically significant ($P=0.09$).

In mice, the body mass index (BMI) can be an indication of metabolic syndrome. We determined the BMI of the mice as follows: (body weight (g)/[nose-anus length (mm)]²). As shown in FIGS. 11A and 11B, *L. reuteri*-derived IL-22 reduces the change in BMI over the course of seven weeks (T7-T0) and six weeks (T7-T1), respectively.

Following seven weeks treatment (T7), animals were killed, tissues were harvested, and liver weights were determined. Liver weights are shown as absolute liver weights (FIG. 12A) or liver weight relative to mouse body weight (FIG. 12B). Both metrics showed that IL-22-secreting *L. reuteri* yielded significantly lower liver weights compared to the wild-type *L. reuteri* or PBS control.

These results show that oral administration of recombinant *L. reuteri* engineered to secrete IL-22 systemically delivers IL-22 in a manner that results in systemic physiological effects. In the present case, the systemic physiological effects included an increase in growth, an increase in growth hormone in the plasma, a reduction in BMI, and a reduction in liver weight. The administration reversed many effects associated with metabolic syndrome. We predict that systemic delivery of IL-22 will also reverse many metabolic symptoms of diabetic subjects, including hyperglycemia and insulin resistance, and will improve insulin sensitivity, preserve gut mucosal barrier and endocrine functions, decrease

endotoxemia and chronic inflammation, and reverse the dysregulation of lipid metabolism in liver and adipose tissues.

Delivery of Peptides Other than IL-22.

L. reuteri can be used to systemically deliver polypeptides other than IL-22. This can be performed by replacing the mIL-22 reading frame from the pVPL3461 plasmid and replacing it with the reading frame of any polypeptide of interest. The edited plasmid can then be introduced into *L. reuteri* using methods described above, and the *L. reuteri* harboring the edited plasmid can be administered as described above. We predict that the *L. reuteri* so modified will be capable of systemically delivering any polypeptide of interest without the bacterium itself being distributed systemically. We predict that diseases and conditions that are alleviated by systemic administration of such polypeptides will be alleviated with the *L. reuteri*-dependent systemic delivery of the peptides.

Statistical Analysis.

In the present examples, ANOVA (analysis of variance) was used for data analysis, and significance in comparisons between groups was analyzed by t-test. Significant difference was considered when P-value is lower than 0.05.

REFERENCES

- Ahrné, S., Molin, G., & Axelsson, L. (1992). Transformation of *Lactobacillus reuteri* with electroporation: Studies on the erythromycin resistance plasmid pLUL631. *Current Microbiology*, 24: 199-205.
- Alvarez-Sieiro P, Montalbin-López M, Mu D, Kuipers O P. Bacteriocins of lactic acid bacteria: extending the family. *Appl Microbiol Biotechnol*. 2016. 100(7):2939-51.
- Bahey-El-Din M, Gahan C G M, Griffin B T. 2010. *Lactococcus lactis* as a cell factory for delivery of therapeutic proteins. *Curr Gene Ther* 10:34-45.
- Barrangou R, van Pijkeren J P. Exploiting CRISPR-Cas immune systems for genome editing in bacteria. *Curr Opin Biotechnol*. 2016 February; 37:61-8.
- Becker S C, Dong S, Baker J R, Foster-Frey J, Pritchard D G, Donovan D M. 2009. LysK CHAP endopeptidase domain is required for lysis of live staphylococcal cells. *FEMS Microbiol Lett* 294:52-60.
- Beisel C L, Gomaa A A, Barrangou R. 2014. A CRISPR design for next-generation antimicrobials. *Genome Biol* 15:516.
- Bikard D, Euler C W, Jiang W, Nussenzweig P M, Goldberg G W, Duportet X, Fischetti V A, Marraffini L A. 2014. Exploiting CRISPR-Cas nucleases to produce sequence-specific antimicrobials. *Nat Biotech* 32:1146-1150.
- Borysowski J, Weber-Dabrowska B, Górski A. Bacteriophage endolysins as a novel class of antibacterial agents. *Exp Biol Med* (Maywood). 2006 April; 231(4):366-77.
- Britton, R. A.; Irwin, R.; Quach, D.; Schaefer, L.; Zhang, J.; Lee, T.; Parameswaran, N.; McCabe, L. R. Probiotic *L. reuteri* Treatment Prevents Bone Loss in a Menopausal Ovariectomized Mouse Model. *J Cell Physiol* 2014, 229 (11), n/a-n/a DOI: 10.1002/jcp.24636.
- Chatel J-M, Pothelune L, Ah-Leung S, Corthier G, Wal J-M, Langella P. 2008. In vivo transfer of plasmid from food-grade transiting lactococci to murine epithelial cells. *Gene Ther* 15:1184-1190.
- Cheng X, Zhang X, Pflugrath J W, Studier F W. 1994. The structure of bacteriophage T7 lysozyme, a zinc amidase and an inhibitor of T7 RNA polymerase. *Proc Natl Acad Sci USA* 91:4034-4038.

- Citorik R J, Mimee M, Lu T K. 2014. Sequence-specific antimicrobials using efficiently delivered RNA-guided nucleases. *Nat Biotech* 32:1141-1145.
- Cotter P D, Ross R P, Hill C. 2013. Bacteriocins [mdash] a viable alternative to antibiotics? *Nat Rev Microbiol* 11:95-105.
- Cronin, M.; Akin, A. R.; Collins, S. A.; Meganck, J.; Kim, J.-B.; Baban, C. K.; Joyce, S. A.; van Dam, G. M.; Zhang, N.; van Sinderen, D.; et al. High resolution in vivo bioluminescent imaging for the study of bacterial tumour targeting. *PLoS ONE* 2012, 7 (1), e30940 DOI: 10.1371/journal.pone.0030940.
- Cronin M, Morrissey D, Rajendran S, El Mashad S M, van Sinderen D, O'Sullivan G C, Tangney M. Orally administered bifidobacteria as vehicles for delivery of agents to systemic tumors. *Mol Ther*. 2010 July; 18(7):1397-407.
- de Azevedo M, Karczewski J, Lefevre F, Azevedo V, Miyoshi A, Wells J M, Langella P, Chatel J-M. 2012. In vitro and in vivo characterization of DNA delivery using recombinant *Lactococcus lactis* expressing a mutated form of *L. monocytogenes* Internalin A. *BMC Microbiol* 12:299.
- de Ruyter, P. G.; Kuipers, O. P.; de Vos, W. M. Controlled gene expression systems for *Lactococcus lactis* with the food-grade inducer nisin. *Appl Environ Microbiol* 1996, 62 (10), 3662-3667.
- De Weirdt R, Crabbé A, Roos S, Vollenweider S, Lacroix C, van Pijkeren J-P, Britton R A, Sarker S, Van de Wiele T, Nickerson C A. 2012. Glycerol Supplementation Enhances *L. reuteri*'s Protective Effect against *S. Typhimurium* Colonization in a 3-D Model of Colonic Epithelium. *PLoS ONE* 7:e37116.
- Dishisha T, Pereyra L P, Pyo S-H, Britton R A, Hatti-Kaul R. 2014. Flux analysis of the *Lactobacillus reuteri* propanediol-utilization pathway for production of 3-hydroxypropionaldehyde, 3-hydroxypropionic acid and 1,3-propanediol from glycerol. *Microb Cell Fact* 13:76.
- Doleyres Y, Beck P, Vollenweider S, Lacroix C. 2005. Production of 3-hydroxypropionaldehyde using a two-step process with *Lactobacillus reuteri*. *Appl Microbiol Biotechnol* 68:467-474.
- Eaton, K. A.; Honkala, A.; Auchtung, T. A.; Britton, R. A. Probiotic *Lactobacillus reuteri* Ameliorates Disease Due to Enterohemorrhagic *Escherichia coli* in Germfree Mice. *Infect Immun* 2011, 79 (1), 185-191 DOI: 10.1128/IAI.00880-10.
- Elzagheid, A.; Algars, A.; Bendardaf, R.; Lamlum, H.; Ristamaki, R.; Collan, Y.; Syrjanen, K.; Pyrhonen, S. E-cadherin expression pattern in primary colorectal carcinomas and their metastases reflects disease outcome. *World J Gastroenterol* 2006, 12 (27), 4304-4309.
- Feliza A, Bourguet, Brian E, Souza, Angela K, Hinz, Matthew A, Coleman, and Paul J, Jackson. Characterization of a Novel Lytic Protein Encoded by the *Bacillus cereus* E33L Gene ampD as a *Bacillus anthracis* Antimicrobial Protein. *Appl Environ Microbiol*. 2012 April; 78(8): 3025-3027.
- Field D, Connor P M O, Cotter P D, Hill C, Ross R P. 2008. The generation of nisin variants with enhanced activity against specific gram-positive pathogens.—PubMed—NCBI. *Mol Microbiol* 69:218-230.
- Field D, Begley M, O'Connor P M, Daly K M, Hugenholtz F, Cotter P D, Hill C, Ross R P. 2012. Bioengineered Nisin A Derivatives with Enhanced Activity against Both Gram Positive and Gram Negative Pathogens. *PLoS ONE* 7:e46884

- Fischetti V A. 2004. The use of phage lytic enzymes to control bacterial infections, p 321-334 In Kutter E, Sulakvelidze A, editors. (ed), Bacteriophages: biology and applications. CRC Press, Boca Raton, Fla.
- Fischetti V A. 2009. Bacteriophage Lysins: the Ultimate Enzybiotic Enzybiotics. John Wiley & Sons, Inc., Hoboken, N.J., USA
- Fogel, M. R.; Gray, G. M. Starch hydrolysis in man: an intraluminal process not requiring membrane digestion. *J Appl Physiol* 1973, 35 (2), 263-267.
- Foster, P. L. Methods for determining spontaneous mutation rates. *Meth Enzymol* 2006, 409, 195-213.
- Frese S A, Benson A K, Tannock G W, Loach D M, Kim J, Zhang M, Oh P L, Heng N C, Patil P B, Juge N, Mackenzie D A, Pearson B M, Lapidus A, Dalin E, Tice H, Goltsman E, Land M, Hauser L, Ivanova N, Kyrpides N C, Walter J. The evolution of host specialization in the vertebrate gut symbiont *Lactobacillus reuteri*. *PLoS Genet*. 2011 February; 7(2):e1001314.
- Gibson D G, Young L, Chuang R-Y, Venter J C, Hutchison C A, Smith H O. 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Meth* 6:343-345.
- Gomaa A A, Klumpe H E, Luo M L, Selle K, Barrangou R, Beisel C L. 2014. Programmable Removal of Bacterial Strains by Use of Genome-Targeting CRISPR-Cas Systems. *mBio* 5:e00928-13.
- Green et al., *Molecular Cloning: A Laboratory Manual*, 4th ed., Cold Spring Harbor Laboratory Press, 2012.
- Guimarães, V. D.; Gabriel, J. E.; Lefevre, F.; Cabanes, D.; Gruss, A.; Cossart, P.; Azevedo, V.; Langella, P. Internalin-expressing *Lactococcus lactis* is able to invade small intestine of guinea pigs and deliver DNA into mammalian epithelial cells. *Microbes Infect* 2005, 7 (5-6), 836-844 DOI: 10.1016/j.micinf.2005.02.012.
- Guimarães V, Innocentin S, Chatel J-M, Lefevre F, Langella P, Azevedo V, 582 Miyoshi A. 2009. A new plasmid vector for DNA delivery using lactococci. *Genet Vaccines Ther* 7:4.
- Huyghebaert N, Vermeire A, Neiryck S, Steidler L, Remaut E, Remon J P. Development of an enteric-coated formulation containing freeze-dried, viable recombinant *Lactococcus lactis* for the ileal mucosal delivery of human interleukin-10. *Eur J Pharm Biopharm*. 2005 August; 60(3):349-59.
- Jacobs C, Huang L J, Bartowsky E, Normark S, Park J T. Bacterial cell wall recycling provides cytosolic muropeptides as effectors for beta-lactamase induction. *EMBO J*. 1994 Oct. 3; 13(19):4684-94.
- Jacobs C, Joris B, Jamin M, Klarsov K, Van Beeumen J, Mengin-Lecreulx D, van Heijenoort J, Park J T, Normark S, Fèbre J M. AmpD, essential for both beta-lactamase regulation and cell wall recycling, is a novel cytosolic N-acetylmuramyl-L-alanine amidase. *Mol Microbiol*. 1995 February; 15(3):553-9.
- Jensen H, Roos S, Jonsson H, Rud I, Grimmer S, van Pijkeren J P, Britton R A, Axelsson L. Role of *Lactobacillus reuteri* cell and mucus-binding protein A (CmbA) in adhesion to intestinal epithelial cells and mucus in vitro. *Microbiology*. 2014 April; 160(Pt 4):671-81.
- Kok, S. D., Stanton, L. H., Slaby, T., Durot, M., Holmes, V. F., Patel, K. G., Platt, D., Shapland, E. B. and Chandran, S. S. (2014). Rapid and reliable DNA assembly via ligase cycling reaction. *ACS synthetic biology*, 3: 97-106.
- Kommineni S, Bretl D J, Lam V, Chakraborty R, Hayward M, Simpson P, Cao Y, Bousounis P, Kristich C J, Salzman

- N H. 2015. Bacteriocin production augments niche competition by enterococci in the mammalian GI tract. *Nature* 526:719-722.
- Lecuit, M.; Ohayon, H.; Braun, L.; Mengaud, J.; Cossart, P. Internalin of *Listeria monocytogenes* with an intact leucine-rich repeat region is sufficient to promote internalization. *Infect Immun* 1997, 65 (12), 5309-5319.
- Liu, Y.; Fatheree, N. Y.; Mangalat, N.; Rhoads, J. M. Human-derived probiotic *Lactobacillus reuteri* strains differentially reduce intestinal inflammation. *Am J Physiol Gastrointest Liver Physiol* 2010, 299 (5), G1087-G1096 DOI: 10.1152/ajpgi.00124.2010.
- Loessner M J. Bacteriophage endolysins—current state of research and applications. *Curr Opin Microbiol*. 2005 August; 8(4):480-7.
- Loessner M J, Kramer K, Ebel F, Scherer S. C-terminal domains of *Listeria monocytogenes* bacteriophage murein hydrolases determine specific recognition and high-affinity binding to bacterial cell wall carbohydrates. *Mol Microbiol*. 2002 April; 44(2):335-49.
- Loonen, L. M., Stolte, E. H., Jaklofsky, M. T., Meijerink, M., Dekker, J., van Baarlen, P., & Wells, J. M. (2014). REG3γ-deficient mice have altered mucus distribution and increased mucosal inflammatory responses to the microbiota and enteric pathogens in the ileum. *Mucosal immunology*, 7: 939-947.
- López R, García E, Garcia P, Garcia J L. The pneumococcal cell wall degrading enzymes: a modular design to create new lysins? *Microb Drug Resist*. 1997; 3(2):199-211.
- Mackenzie D A, Jeffers F, Parker M L, Vibert-Vallet A, Bongaerts R J, Roos S, Walter J, Juge N. Strain-specific diversity of mucus-binding proteins in the adhesion and aggregation properties of *Lactobacillus reuteri*. *Microbiology*. 2010 November; 156(Pt 11):3368-78. Oh J H, van Pijkeren J P. CRISPR-Cas9-assisted recombineering in *Lactobacillus reuteri*. *Nucleic Acids Res*. 2014; 42(17): e131.
- Navarre W W, Ton-That H, Faull K F, Schneewind O. 1999. Multiple Enzymatic Activities of the Murein Hydrolase from Staphylococcal Phage φ11. *Journal of Biological Chemistry* 274:15847-15856.
- Nelson D, Schuch R, Chahales P, Zhu S, Fischetti V A. 2006. PlyC: A multimeric bacteriophage lysin. *Proc Natl Acad Sci USA* 103:10765-10770.
- Nicoletti M, Bertani G. 1983. DNA fusion product of phage P2 with plasmid pBR322: A new phasmid. *Mol Gen Genet* 189:343-347.
- Oh, P. L.; Benson, A. K.; Peterson, D. A.; Patil, P. B.; Moriyama, E. N.; Roos, S.; Walter, J. Diversification of the gut symbiont *Lactobacillus reuteri* as a result of host-driven evolution. *ISME J* 2010, 4 (3), 377-387 DOI: 10.1038/ismej.2009.123.
- Oh J H, van Pijkeren J P. CRISPR-Cas9-assisted recombineering in *Lactobacillus reuteri*. *Nucleic Acids Res*. 2014; 42(17):e131.
- Paez-Espino D, Morovic W, Sun C L, Thomas B C, Ueda K-I, Stahl B, Barrangou R, Banfield J F. 2013. Strong bias in the bacterial CRISPR elements that confer immunity to phage. *Nat Comms* 4:1430-1430.
- Pálffy, R.; Gardlik, R.; Hodosy, J.; Behuliak, M.; Resko, P.; Radvinsky, J.; Celec, P. Bacteria in gene therapy: bacteriofection versus alternative gene therapy. *Gene Ther* 2006, 13 (2), 101-105 DOI:10.1038/sj.gt.3302635.
- Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic acids research*, 29: e45-e45.

- Pfaffl, M. W., Horgan, G. W., & Dempfle, L. (2002). Relative expression software tool (REST©) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic acids research*, 30: e36-e36.
- Rajpal G, Liu M, Zhang Y, Arvan P. Single-chain insulins as receptor agonists. *Mol Endocrinol*. 2009 May; 23(5):679-88.
- Riboulet-Bisson E, Sturme M H J, Jeffery I B, O'Donnell M M, Neville B A, Forde B M, Claesson M J, Harris H, Gardiner G E, Casey P G, Lawlor P G, O'Toole P W, Ross R P. 2012. Effect of *Lactobacillus salivarius* Bacteriocin Abp118 on the Mouse and Pig Intestinal Microbiota. *PLoS ONE* 7:e31113.
- Riedel, C. U.; Monk, I. R.; Casey, P. G.; Morrissey, D.; O'Sullivan, G. C.; Tangney, M.; Hill, C.; Gahan, C. G. M. Improved luciferase tagging system for *Listeria monocytogenes* allows real-time monitoring in vivo and in vitro. *Appl Environ Microbiol* 2007, 73 (9), 3091-3094 DOI: 10.1128/AEM.02940-06.
- Robert S, Steidler L. 2014. Recombinant *Lactococcus lactis* can make the difference in antigen-specific immune tolerance induction, the Type 1 Diabetes case. *Microb Cell Fact* 13:S11.
- Rosche, W. A.; Foster, P. L. Determining mutation rates in bacterial populations. *Methods* 2000, 20 (1), 4-17.
- Sabat R, Ouyang W, Wolk K. Therapeutic opportunities of the IL-22-IL-22R1 system. *Nat Rev Drug Discov*. 2014 January; 13(1):21-38.
- Schaefer L, Auchtung T A, Hermans K E, Whitehead D, Borhan B, Britton R A. 2010. The antimicrobial compound reuterin (3-hydroxypropionaldehyde) induces oxidative stress via interaction with thiol groups. *Microbiology* 156:1589-1599.
- Sheehan M M, Stanley E, Fitzgerald G F, van Sinderen D. 1999. Identification and characterization of a lysis module present in a large proportion of bacteriophages infecting *Streptococcus thermophilus*. *Appl Environ Microbiol* 65:569-577.
- Shi, Y.; Yan, Y.; Ji, W.; Bin Du; Meng, X.; Wang, H.; Sun, J. Characterization and determination of holin protein of *Streptococcus suis* bacteriophage SMP in heterologous host. *Virol J* 2012, 9 (1), 70-70 DOI: 10.1186/1743-422X-9-70.
- Singh K, Kadesjö E, Lindroos J, Hjort M, Lundberg M, Espes D, Carlsson P O, Sandler S, Thorvaldson L. Interleukin-35 administration counteracts established murine type 1 diabetes—possible involvement of regulatory T cells. *Sci Rep*. 2015 Jul. 30; 5:12633.
- Sovran, B., Loonen, L. M., Lu, P., Hugenholtz, F., Belzer, C., Stolte, E. H., . . . & Dekker, J. (2015). IL-22-STAT3 pathway plays a key role in the maintenance of ileal homeostasis in mice lacking secreted mucus barrier. *Inflammatory bowel diseases*, 21: 531-542.
- Spinler J K, Taweechoatipatr M, Rognerud C L, Ou C N, Tumwasorn S, Versalovic J. 2008. Human-derived probiotic *Lactobacillus reuteri* demonstrate antimicrobial activities targeting diverse enteric bacterial pathogens. *Anaerobe* 14:166-171.
- Steidler L, Hans W, Schotte L, Neiryneck S, Obermeier F, Falk W, Fiers W, Remaut E. Treatment of murine colitis by *Lactococcus lactis* secreting interleukin-10. *Science*. 2000 Aug. 25; 289(5483): 1352-5.
- Sulakvelidze, Alexander; Alavidze, Zemphira; and J. Glenn Morris, Jr. Bacteriophage Therapy. *Antimicrob Agents Chemother*. 2001, 45(3): 649-659.

- Summers W C. Bacteriophage therapy. *Annu Rev Microbiol*. 2001; 55:437-51.
- Sun J, Furio L, Mecheri R, van der Does A M, Lundeberg E, Saveanu L, Chen Y, van Endert P, Agerberth B, Diana J. Pancreatic β -Cells Limit Autoimmune Diabetes via an Immunoregulatory Antimicrobial Peptide Expressed under the Influence of the Gut Microbiota. *Immunity*. 2015 Aug. 18; 43(2):304-17.
- Sznol, M.; Lin, S. L.; Bermudes, D.; Zheng, L.-M.; King, I. Use of preferentially replicating bacteria for the treatment of cancer. *Journal of Clinical Investigation* 2000, 105 (8), 1027-1030 DOI:10.1172/JCI9818.
- Talarico T L, Casas I A, Chung T C, Dobrogosz W J. 1988. Production and isolation of reuterin, a growth inhibitor produced by *Lactobacillus reuteri*. *Antimicrob Agents Chemother* 32:1854-1858.
- Tannock, G. W.; Wilson, C. M.; Loach, D.; Cook, G. M.; Eason, J.; O'Toole, P. W.; Holtrop, G.; Lawley, B. Resource partitioning in relation to cohabitation of *Lactobacillus* species in the mouse forestomach. *ISME J* 2011, 6 (5), 927-938 DOI: 10.1038/ismej.2011.161.
- Thomas, C. M.; Hong, T.; van Pijkeren, J.P.; Hemarajata, P.; Trinh, D. V.; Hu, W.; Britton, R. A.; Kalkum, M.; Versalovic, J. Histamine Derived from Probiotic *Lactobacillus reuteri* Suppresses TNF via Modulation of PKA and ERK Signaling. *PLoS ONE* 2012, 7 (2), e31951 DOI: 10.1371/journal.pone.0037116.g006.
- van Pijkeren, J.P.; Morrissey, D.; Monk, I. R.; Cronin, M.; Rajendran, S.; O'Sullivan, G. C.; Gahan, C. G. M.; Tangney, M. A novel *Listeria monocytogenes*-based DNA delivery system for cancer gene therapy. *Hum. Gene Ther*. 2010, 21 (4), 405-416 DOI: 10.1089/hum.2009.022.
- van Pijkeren J P, Britton R A. High efficiency recombineering in lactic acid bacteria. *Nucleic Acids Res*. 2012 May; 40(10):e76.
- van Pijkeren, J.P.; Britton, R. A. Precision genome engineering in lactic acid bacteria. *Microb Cell Fact* 2014, 13 Suppl 1, S10-S10 DOI: 10.1186/1475-2859-13-S1-S10.
- van Pijkeren J-P, Neoh K M, Sirias D, Findley A S, Britton R A. 2012. Exploring optimization parameters to increase ssDNA recombineering in *Lactococcus lactis* and *Lactobacillus reuteri*. *Bioengineered* 3:209-217.
- Wang I N, Smith D L, Young R. 2000. Holins: the protein clocks of bacteriophage infections. *Annual Reviews in Microbiology*.
- Wang X, Ota N, Manzanillo P, Kates L, Zavala-Solorio J, Eidenschenk C, Zhang J, Lesch J, Lee W P, Ross J, Diehl L, van Bruggen N, Kolumam G, Ouyang W. Interleukin-22 alleviates metabolic disorders and restores mucosal immunity in diabetes. *Nature*. 2014 Oct. 9; 514(7521): 237-41.
- Wells M. et al. *Lactococcus lactis*: high-level expression of tetanus toxin fragment C and protection against lethal challenge. *Mol. Microbiol.*, 8 (1993), pp. 1155-1162.
- Zhang Y, Eigenbrot C, Zhou L, Shia S, Li W, Quan C, Tom J, Moran P, Di Lello P, Skelton N J, Kong-Beltran M, Peterson A, Kirchhofer D. Identification of a small peptide that inhibits PCSK9 protein binding to the low density lipoprotein receptor. *J Biol Chem*. 2014 Jan. 10; 289(2):942-55.
- Zhou, Y; Liang, Y; Lynch, K. H.; Dennis, J. J.; Wishart, D. S. PHAST: A Fast Phage Search Tool. 2011, 39 (suppl), W347-W352 DOI: 10.1093/nar/gkr485.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 32

<210> SEQ ID NO 1
 <211> LENGTH: 311
 <212> TYPE: DNA
 <213> ORGANISM: Lactobacillus reuteri

<400> SEQUENCE: 1

```
taccaagaat aactttcatc gtaaaaggca agtaattgag gaaacttgaa gtttttctct    60
attacttgcc ttctttattht tattaagcta aatatgtttt aaataattaa ctataacgga    120
cctgcttggc ggaaactaaa cagtaagaac tttaaattat aaaaatctgc aaccgttttc    180
taaaatthtg cgcaagcggg tgcgcaaaat ttttaaattt gatattatta atattgcaat    240
aattcatgaa gcgcttataa taatcacaag tgtcttttag aactatttta taagttaagg    300
agttgttagc a                                                    311
```

<210> SEQ ID NO 2
 <211> LENGTH: 146
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```
Ala Pro Ile Ser Ser His Cys Arg Leu Asp Lys Ser Asn Phe Gln Gln
1           5           10           15
Pro Tyr Ile Thr Asn Arg Thr Phe Met Leu Ala Lys Glu Ala Ser Leu
20           25           30
Ala Asp Asn Asn Thr Asp Val Arg Leu Ile Gly Glu Lys Leu Phe His
35           40           45
Gly Val Ser Met Ser Glu Arg Cys Tyr Leu Met Lys Gln Val Leu Asn
50           55           60
Phe Thr Leu Glu Glu Val Leu Phe Pro Gln Ser Asp Arg Phe Gln Pro
65           70           75           80
Tyr Met Gln Glu Val Val Pro Phe Leu Ala Arg Leu Ser Asn Arg Leu
85           90           95
Ser Thr Cys His Ile Glu Gly Asp Asp Leu His Ile Gln Arg Asn Val
100          105          110
Gln Lys Leu Lys Asp Thr Val Lys Lys Leu Gly Glu Ser Gly Glu Ile
115          120          125
Lys Ala Ile Gly Glu Leu Asp Leu Leu Phe Met Ser Leu Arg Asn Ala
130          135          140
Cys Ile
145
```

<210> SEQ ID NO 3
 <211> LENGTH: 209
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Recombinant IL-35

<400> SEQUENCE: 3

```
Arg Lys Gly Pro Pro Ala Ala Leu Thr Leu Pro Arg Val Gln Cys Arg
1           5           10           15
Ala Ser Arg Tyr Pro Ile Ala Val Asp Cys Ser Trp Thr Leu Pro Pro
20           25           30
Ala Pro Asn Ser Thr Ser Pro Val Ser Phe Ile Ala Thr Tyr Arg Leu
35           40           45
```


-continued

Gly Met Ala Ala Arg Gly His Ser Trp Pro Cys Leu Gln Gln Thr Pro
 50 55 60

Thr Ser Thr Ser Cys Thr Ile Thr Asp Val Gln Leu Phe Ser Met Ala
 65 70 75 80

Pro Tyr Val Leu Asn Val Thr Ala Val His Pro Trp Gly Ser Ser Ser
 85 90 95

Ser Phe Val Pro Phe Ile Thr Glu His Ile Ile Lys Pro Asp Pro Pro
 100 105 110

Glu Gly Val Arg Leu Ser Pro Leu Ala Glu Arg Gln Leu Gln Val Gln
 115 120 125

Trp Glu Pro Pro Gly Ser Trp Pro Phe Pro Glu Ile Phe Ser Leu Lys
 130 135 140

Tyr Trp Ile Arg Tyr Lys Arg Gln Gly Ala Ala Arg Phe His Arg Val
 145 150 155 160

Gly Pro Ile Glu Ala Thr Ser Phe Ile Leu Arg Ala Val Arg Pro Arg
 165 170 175

Ala Arg Tyr Tyr Val Gln Val Ala Ala Gln Asp Leu Thr Asp Tyr Gly
 180 185 190

Glu Leu Ser Asp Trp Ser Leu Pro Ala Thr Ala Thr Met Ser Leu Gly
 195 200 205

Lys

<210> SEQ ID NO 4
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Peptide Linker

<400> SEQUENCE: 4

Gln Arg Gly Gly Gly Gly Gln Arg
 1 5

<210> SEQ ID NO 5
 <211> LENGTH: 146
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Val Pro Ile Gln Lys Val Gln Asp Asp Thr Lys Thr Leu Ile Lys Thr
 1 5 10 15

Ile Val Thr Arg Ile Asn Asp Ile Ser His Thr Gln Ser Val Ser Ser
 20 25 30

Lys Gln Lys Val Thr Gly Leu Asp Phe Ile Pro Gly Leu His Pro Ile
 35 40 45

Leu Thr Leu Ser Lys Met Asp Gln Thr Leu Ala Val Tyr Gln Gln Ile
 50 55 60

Leu Thr Ser Met Pro Ser Arg Asn Val Ile Gln Ile Ser Asn Asp Leu
 65 70 75 80

Glu Asn Leu Arg Asp Leu Leu His Val Leu Ala Phe Ser Lys Ser Cys
 85 90 95

His Leu Pro Trp Ala Ser Gly Leu Glu Thr Leu Asp Ser Leu Gly Gly
 100 105 110

Val Leu Glu Ala Ser Gly Tyr Ser Thr Glu Val Val Ala Leu Ser Arg
 115 120 125

Leu Gln Gly Ser Leu Gln Asp Met Leu Trp Gln Leu Asp Leu Ser Pro
 130 135 140

-continued

Gly Cys
145

<210> SEQ ID NO 6
<211> LENGTH: 39
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Phe Ala Leu Leu Gly Asp Phe Phe Arg Lys Ser Lys Glu Lys Ile Gly
1 5 10 15

Lys Glu Phe Lys Arg Ile Val Gln Arg Ile Lys Asp Phe Leu Arg Asn
20 25 30

Leu Val Pro Arg Thr Glu Ser
35

<210> SEQ ID NO 7
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Peptide PCSK9 Inhibitor

<400> SEQUENCE: 7

Thr Val Phe Thr Ser Trp Glu Glu Tyr Leu Asp Trp Val
1 5 10

<210> SEQ ID NO 8
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer oVPL329

<400> SEQUENCE: 8

attccttgga cttcatttac tgggtttaac 30

<210> SEQ ID NO 9
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer oVPL363

<400> SEQUENCE: 9

taatagaga taatgccgac tgtac 25

<210> SEQ ID NO 10
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer oVPL1219

<400> SEQUENCE: 10

ttcatgggga tgaatgcttc tgctaataca ttaccagtta atactcgttg 50

<210> SEQ ID NO 11
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer oVPL1220

<400> SEQUENCE: 11

-continued

cttggttttc taattttggt tcaaagatca aacacaagca ttacgtaaac tc 52

<210> SEQ ID NO 12
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1221

<400> SEQUENCE: 12

gcttgaaacg ttcaattgaa atggca 26

<210> SEQ ID NO 13
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1222

<400> SEQUENCE: 13

tgtaaaacca ataaggactg aagc 24

<210> SEQ ID NO 14
 <211> LENGTH: 60
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1223

<400> SEQUENCE: 14

ggagttgctt cagtccttat tggttttaca ttcattgggga tgaatgcttc tgctaataca 60

<210> SEQ ID NO 15
 <211> LENGTH: 60
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1224

<400> SEQUENCE: 15

tgatctttga accaaaatta gaaaaccaag gcttgaaacg ttcaattgaa atggcaatta 60

<210> SEQ ID NO 16
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1313

<400> SEQUENCE: 16

actccctgaa gaatataccc tcc 23

<210> SEQ ID NO 17
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1314

<400> SEQUENCE: 17

cgctattgag cacagatag ag 22

<210> SEQ ID NO 18
 <211> LENGTH: 22
 <212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1315

 <400> SEQUENCE: 18

 atgcttcccc gtataaccat ca 22

<210> SEQ ID NO 19
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1316

 <400> SEQUENCE: 19

 ggccatatct gcatcatacc ag 22

<210> SEQ ID NO 20
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1321

 <400> SEQUENCE: 20

 gatcacccgac aagggcctg 19

<210> SEQ ID NO 21
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1322

 <400> SEQUENCE: 21

 ggctatgaaa ctcgactgc c 21

<210> SEQ ID NO 22
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1325

 <400> SEQUENCE: 22

 ggctgtattc ccctccatcg 20

<210> SEQ ID NO 23
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer oVPL1326

 <400> SEQUENCE: 23

 ccagttggta acaatgccat gt 22

<210> SEQ ID NO 24
 <211> LENGTH: 500
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: gVPL1 Codon-Optimized IL-22 Coding Sequence

 <400> SEQUENCE: 24

 attcatgggg atgaatgctt ctgctaatac attaccagtt aatactcgtt gtaaattaga 60

-continued

```

agttagtaat tttcaacaac catatattgt taatcgact tttatgtag ctaaagaagc 120
tagtttagct gataataata ctgatgttcg ttaattggg gaaaaattat ttcgtggtgt 180
tagtgctaaa gatcaatggt atttaatgaa acaagtttta aattttactt tagaagatgt 240
tttattacca caaagtgatc gttttcaacc atatatgcaa gaagttgttc catttttaac 300
taaattaagt aatcaattaa gtagttgtca tattagtggg gatgatcaaa atattcaaaa 360
aaatgttcgt cgtttaaaag aaactgttaa aaaattaggg gaaagtgggt aaattaaagc 420
tattggtgaa ttagatttat ttttatgag tttacgtaat gcttgtgttt gatctttgaa 480
ccaaaattag aaaaccaagg 500

```

<210> SEQ ID NO 25

<211> LENGTH: 5090

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Vector pVPL3461

<400> SEQUENCE: 25

```

aatgattgaa aacaaaattg actttaatga aaaaagaaat ttgatgattt catcggttat 60
tttagtaatt gggattggaa atgcctatct tcaattagga aaatatcaat tctctggatt 120
agctgttgca gctgtcctcg ggataataat gaatttaatc cttccacaaa aagccaaaag 180
tgaaatgtag tttgaatttt taagcagagt aagaaaaggc atccgacctc ttactctgct 240
tttttcgtat gataacaaaa caaaaaatta agaggctcgtg gaaaaaacag attgtaagaa 300
aaagtttatc aagtctaaga aaacgttttc ttacactct tatcaattga ttagtaggta 360
tatataaacc taattttttt tgaattttta cgaaattaat gatataataa cattcgtaac 420
ttggggtagt tcgttttatt tgcaacgttc atccaattta caagccaaat tattgttata 480
aggaaagagg tggaatatga atgttatcga agaataatcg aaaggaacaa ttccggaaac 540
aagagccgaa aaagcaacgt tttgcaatta aaaagctcac tgtcggagtt gcttcagtcc 600
ttattggttt tacattcatg gggatgaatg cttctgctaa tacattacca gtttaactc 660
gttgtaaatt agaagttagt aattttcaac aaccatatat tgtaaatcgt acttttatgt 720
tagctaaaga agctagttta gctgataata atactgatgt tcgtttaatt ggtgaaaaat 780
tatttcgtgg tgtagtgct aaagatcaat gttatttaat gaaacaagtt ttaaatttta 840
ctttagaaga tgttttatta ccacaaagtg atcgttttca accatatatg caagaagttg 900
ttccattttt aactaaatta agtaatcaat taagtagttg tcatattagt ggtgatgatc 960
aaaatattca aaaaaatggt cgctgtttta aagaaactgt taaaaatta ggtgaaagtg 1020
gtgaaattaa agctattggt gaattagatt tattatttat gagtttacgt aatgcttggt 1080
ttgatcttt gaacccaaat tagaaaacca aggtctctgc taatacattg aaacgttcaa 1140
ttgaaatggc aattaaacaa attacagcac gtgttgcttt gattgatagc caaaaagcag 1200
cagttgataa agcaattact gatattgctg aaaaattgta atttataaat aaaaatcacc 1260
ttttagaggg ggttttttta tttataaatt attcgtttga tttcgctttc gatagaacaa 1320
tcaaagcgag attataaaag ccagtcatta ggctatctg acaattcctg aatagagttc 1380
ataaacaatc cagcatgata accatcacia acagaatgat gtacctgtaa agatagcggg 1440
aaatatattg aattaccttt attaataaat tttcctgctg taataatggg tagaaggtaa 1500
ttactattat tattgatatt taagttaaac ccagtaaatg aagtccaagg aataatagaa 1560

```


-continued

agagaaaaag	cattttcagg	tataggtggt	ttgggaaaca	atttccccga	accattatat	1620
ttctctacat	cagaaaggta	taaatcataa	aactctttga	agtcattctt	tacaggagtc	1680
caaataaccag	agaatgtttt	agatacacca	tcaaaaattg	tataaagtgg	ctctaactta	1740
tccaataaac	ctaactctcc	gtcgctattg	taaccagttc	taaaagctgt	atgtgagttt	1800
atcacccctg	tcactaagaa	aataaatgca	gggtaaaatt	tatatccttc	ttgttttatg	1860
tttcggtata	aaacactaat	atcaatttct	gtggttatac	taaaagtcgt	ttgttggttc	1920
aaataatgat	taaatatctc	ttttctcttc	caattgtcta	aatcaatttt	attaaagtcc	1980
atgggtttca	ctctccttct	acatttttta	acctaataat	gccaaatacc	gtttgccacc	2040
cctctctttt	gataattata	atattggcga	aattcgcttc	taaagatgaa	acgcaatatt	2100
atatgcttgc	tttataattg	tgagcgctca	caatttatgt	gattatacca	gccccctcac	2160
tacatgtcaa	gaataaactg	ccaaagcata	atgtaaggaa	gataaatccc	ataagggcgg	2220
gagcagaatg	tccgagacta	atgtaaattt	gtccaccaat	taaaggaccg	ataacgcgag	2280
cttctcgagt	gcatattttc	ggcaatcttc	tcaatgagat	gctcttcagc	atgttcaatg	2340
atgtcgattt	tttattaaaa	cgtctcaaaa	tcgtttctga	gacgttttag	cgtttatttc	2400
gtttagttat	cggcataatc	gttaaacag	gcgttatcgt	agcgtaaaag	cccttgagcg	2460
tagcgtgctt	tgcagcgaag	atgttgtctg	ttagattatg	aaagccgatg	actgaatgaa	2520
ataataagcg	cagcgtcctt	ctatttcggt	tggaggaggc	tcaagggagt	ttgaggggat	2580
gaaattccct	catgggtttg	attttaaaaa	ttgcttgcaa	ttttgccgag	cggtagcgtc	2640
ggaaaatttt	tgaaaaaaat	ttggaatttg	gaaaaaatg	gggggaaagg	aagcgaattt	2700
tgttccgta	ctacgacccc	ccattaagtg	ccgagtgcc	atttttgtgc	caaaaacgct	2760
ctatcccaac	tggctcaagg	gtttgagggg	ttttcaatc	gccaacgaat	cgccaacggt	2820
ttcgccaacg	ttttttataa	atctatattt	aagtagcttt	attggtggtt	ttatgattac	2880
aaagtgatac	actaatttta	taaaattatt	tgattggagt	tttttaaag	gtgatttcag	2940
aatcgaaaaa	aagagttatg	atctctctga	caaaagagca	agataaaaaa	ttaacagata	3000
tggcgaaaca	aaaaggtttt	tcaaaatctg	cggttgccgc	gtagctata	gaagaatatg	3060
caagaaagga	atcagaacaa	aaaaaataag	cgaaagctcg	cgtttttaga	aggatacgag	3120
ttttcgctac	ttgtttttga	taaggtaata	tatcatggct	attaaaaata	ctaaagctag	3180
aaattttggg	tttttattat	atcctgactc	aattccta	gattggaaag	aaaaattaga	3240
gagtttgggc	gtatctatgg	ctgtcagtc	ttacacgat	atggacgaaa	aaaaagataa	3300
agatacatgg	aatagtagtg	atgttatacg	aatggaaag	cactataaaa	aaccacacta	3360
tcacgttata	tatattgcac	gaaatcctgt	aacaatagaa	agcgttagga	acaagattaa	3420
gcgaaaattg	gggaatagtt	cagttgctca	tggtgagata	cttgattata	tcaaaggttc	3480
atatgaatat	ttgactcatg	aatcaaagga	cgctattgct	aagaataaac	atatatacga	3540
caaaaaagat	attttgaaca	ttaatgattt	tgatattgac	cgctatataa	cacttgatga	3600
aagccaaaaa	agagaattga	agaatttact	tttagatata	gtggatgact	ataatttggt	3660
aaatacaaaa	gatttaattg	cttttattcg	ccttagggga	gcgagtttg	gaattttaaa	3720
tacgaatgat	gtaaaagata	ttgtttcaac	aaactctagc	gcctttagat	tatggtttga	3780
gggcaattat	cagtggtgat	atagagcaag	ttatgcaaag	gttcttgatg	ctgaaacggg	3840
ggaaataaaa	tgacaaacaa	agaaaaagag	ttatttgctg	aaaatgagga	attaaaaaaa	3900
gaaattaagg	acttaaaaga	gcgtattgaa	agatacagag	aatggaagt	tgaattaagt	3960

-continued

```

acaacaatag atttattgag aggagggatt attgaataaa taaaagcccc cctgacgaaa 4020
gtcgcaggca atagttacc ttattatcaa gataagaaag aaaaggattt ttcgctacgc 4080
tcaaatcctt taaaaaaca caaaagacca cattttttaa tgtggctctt attcttcaac 4140
taaagcacc attagttcaa caaacgaaaa ttggataaag tgggatattt ttaaaatata 4200
tatttatggt acagtaatat tgacttttaa aaaaggattg attctaataga agaaagcaga 4260
caagtaagcc tcctaaattc actttagata aaaatttagg aggcatatca aatgaacgag 4320
aaaaatataa aacacagtca aaactttatt acttcaaac ataatataga taaaataatg 4380
acaaatataa gattaaatga acatgataat atctttgaaa tcggctcagg aaaaggccat 4440
tttacccttg aattagtaaa gaggtgtaat ttcgtaactg ccattgaaat agaccataaa 4500
ttatgcaaaa ctacagaaaa taaacttggt gatcacgata attccaagt ttaaacaaag 4560
gatataattg agtttaaac tcctaaaaac caatcctata aaatatatgg taatatacct 4620
tataacataa gtacggatat aatagcaaaa attgttttg atagtatagc taatgagatt 4680
tatttaatcg tgaatacgg gtttgctaaa agattattaa atacaaaacg ctcattggca 4740
ttacttttaa tggcagaagt tgatatttct atattaagta tggttccaag agaatatttt 4800
catcctaac ctaaagtgaa tagctcactt atcagattaa gtagaaaaaa atcaagaata 4860
tcacacaaag ataaacaaaa gtataattat ttcgttatga aatggggtta caaagaatac 4920
aagaaaatat ttacaaaaaa tcaatttaac aattccttaa aacatgcagg aattgacgat 4980
ttaaacaata ttagctttga acaattctta tctcttttca atagctataa attatttaat 5040
aagtaataat atgagataat gccgactgta ctttttacag tcggttttct 5090

```

```

<210> SEQ ID NO 26
<211> LENGTH: 500
<212> TYPE: DNA
<213> ORGANISM: Lactobacillus reuteri

```

```

<400> SEQUENCE: 26

```

```

aatgattgaa aacaaaattg actttaatga aaaaagaaat ttgatgattt catcggttat 60
tttagtaatt gggattgaa atgcctatct tcaattagga aaatatcaat tctctggatt 120
agctgttgca gctgtcctcg ggataataat gaatttaatc ctccacaaa aagccaaaag 180
tgaaatgtag tttgaatttt taagcagagt aagaaaaggc atccgacctc ttactctgct 240
tttttcgtat gataacaaaa caaaaaatta agaggctgtg gaaaaaacag attgtaagaa 300
aaagtttatc aagtctaaga aaacgttttc ttactactct tatcaattga ttagtaggta 360
tatataaacc taattttttt tgaattttta cgaaattaat gatataataa cattcgtaac 420
ttggggtagt tcgttttatt tgcaacgttc atccaattta caagccaaat tattgttata 480
aggaaagagg tggaatatga 500

```

```

<210> SEQ ID NO 27
<211> LENGTH: 144
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Artificial signal peptide in pVPL3461

```

```

<400> SEQUENCE: 27

```

```

atgttatcga agaataatcg aaaggaacaa ttccggaaac aagagccgaa aaagcaacgt 60
tttgcaatta aaaagctcac tgcggaggtt gcttcagtcc ttattggttt tacattcatg 120

```


-continued

 gggatgaatg cttctgctaa taca 144

<210> SEQ ID NO 28
 <211> LENGTH: 438
 <212> TYPE: DNA
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 28

ttaccagtta atactcgttg taaattagaa gttagtaatt ttcaacaacc atatattggt 60
 aatcgtagctt ttatgtagc taaagaagct agtttagctg ataataatac tgatgttcgt 120
 ttaattggtg aaaaattatt tcgtgggtgtt agtgctaaag atcaatgta tttaatgaaa 180
 caagttttaa attttacttt agaagatggt ttattaccac aaagtgatcg ttttcaacca 240
 tatatgcaag aagttgttcc atttttaact aaattaagta atcaattaag tagttgtcat 300
 attagtgggtg atgatcaaaa tattcaaaaa aatgttcgtc gtttaaaaga aactgttaaa 360
 aaattaggtg aaagtgggtga aattaaagct attgggtgaat tagatttatt atttatgagt 420
 ttacgtaatg cttgtgtt 438

<210> SEQ ID NO 29
 <211> LENGTH: 70
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Inverted repeat in pVPL3461

<400> SEQUENCE: 29

aaaaattgta atttataaat aaaaatcacc ttttagaggt ggttttttta tttataaatt 60
 attcgtttga 70

<210> SEQ ID NO 30
 <211> LENGTH: 648
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chloramphenicol marker in pVPL3461

<400> SEQUENCE: 30

atgaacttta ataaaattga tttagacaat tggaagagaa aagagatatt taatcattat 60
 ttgaaccaac aaacgacttt tagtataacc acagaaattg atattagtgt tttataccga 120
 aacataaaac aagaaggata taaattttac cctgcattta ttttcttagt gacaaggggtg 180
 ataaaactcaa atacagcttt tagaactggt tacaatagcg acggagagtt aggttattgg 240
 gataagttag agccacttta tacaatTTTT gatggtgtat ctaaaacatt ctctggtatt 300
 tggactcctg taaagaatga cttcaaagag ttttatgatt tataccttcc tgatgtagag 360
 aaatataatg gttcggggaa attgtttccc aaaacaccta tacctgaaaa tgctttttct 420
 ctttctatta ttccttgac ttcatttact gggtttaact taaatatcaa taataatagt 480
 aattaccttc taccattat tacagcagga aaattcatta ataaaggtaa ttcaatatat 540
 ttaccgctat ctttacaggt acatcattct gtttgtgatg gttatcatgc tggattgttt 600
 atgaactcta ttcaggaatt gtcagatagg cctaatgact ggctttta 648

<210> SEQ ID NO 31
 <211> LENGTH: 211
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Phelp promoter in pVPL3461

-continued

<400> SEQUENCE: 31

```

cattatgctt tggcagttta ttcttgacat gtagtgaggg ggctggtata atcacataaa      60
ttgtgagcgc tcacaattat aaagcaagca tataatattg cgtttcatct ttagaagcga      120
atttcgccaa tattataatt atcaaaagag aggggtggca aacggtatth ggcattatta      180
ggtaaaaaaa tgtagaagga gagtgaaacc c                                     211

```

<210> SEQ ID NO 32

<211> LENGTH: 735

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Erythromycin marker in pVPL3461

<400> SEQUENCE: 32

```

atgaacgaga aaaatataaa acacagtcaa aactttatta cttcaaaaca taatatagat      60
aaaataatga caaatataag attaatgaa catgataata tctttgaaat cggctcagga      120
aaaggccatt ttacccttga attagtaaag aggtgtaatt tcgtaactgc cattgaaata      180
gaccataaat tatgcaaaac tacagaaaat aaacttggtg atcacgataa tttccaagtt      240
ttaaacaagg atatattgca gtttaaattt ctaaaaaacc aatectataa aatatatggt      300
aatatacctt ataacataag tacggatata atacgcaaaa ttgtttttga tagtatagct      360
aatgagattt atttaatcgt ggaatacggg tttgctaaaa gattattaaa tacaaaacgc      420
tcattggcat tacttttaat ggcagaagtt gatatttcta tattaagtat ggttccaaga      480
gaatattttc atcctaaacc taaagtgaat agctcactta tcagattaag tagaaaaaaa      540
tcaagaatat cacacaaaga taaacaaaag tataattatt tcgttatgaa atgggttaac      600
aaagaatata agaaaatatt tacaaaaaat caatttaaca attccttaaa acatgcagga      660
attgacgatt taaacaatat tagctttgaa caattcttat ctcttttcaa tagctataaa      720
ttatttaata agtaa                                                    735

```

We claim:

1. A method of systemically introducing a polypeptide into a bloodstream of a subject, the method comprising administering into the gastrointestinal tract of the subject a bacterium that produces and releases the polypeptide, wherein the bacterium comprises a recombinant gene configured to express the polypeptide, wherein the bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream of the subject in a detectable amount without the bacterium being introduced in the bloodstream of the subject in a detectable amount, wherein the bacterium is *Lactobacillus reuteri*.

2. The method of claim 1, wherein the bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream in an amount effective to induce at least one direct systemic effect in the subject.

3. The method of claim 1, wherein the bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream in an amount effective to induce at least one direct effect in a non-gastrointestinal tissue in the subject.

4. The method of claim 1, wherein the bacterium is administered in an amount effective to introduce the polypeptide in the bloodstream in an amount effective to induce at least one direct effect in a tissue selected from the group consisting of liver, muscles, lungs, kidneys, pancreas, and adipose tissue in the subject.

5. The method of claim 1, wherein the subject suffers from a condition treatable with systemic introduction of the polypeptide and wherein the polypeptide is introduced in the bloodstream of the subject in an amount effective to treat the condition.

6. The method of claim 1, wherein the subject suffers from a condition treatable with systemic introduction of the polypeptide but not treatable with local introduction of the polypeptide to the gastrointestinal tract without systemic introduction of the polypeptide, and wherein the polypeptide is introduced in the bloodstream of the subject in an amount effective to treat the condition.

7. The method of claim 1, wherein the polypeptide is selected from the group consisting of a cytokine, a hormone, an antibody, an antimicrobial peptide, and an antigenic peptide.

8. The method of claim 1, wherein the polypeptide is selected from the group consisting of interleukin-22 (IL-22), interleukin-35 (IL-35), insulin, leptin, cathelicidin related antimicrobial peptide, a peptide inhibitor of proprotein convertase subtilisin/kexin type 9 (PCSK9), and an endolysin.

9. The method of claim 8, wherein the subject suffers from at least one condition selected from the group consisting of insulin resistance, hyperglycemia, lipid dysregulation, hyperlipidemia, and obesity, and wherein the polypeptide is

introduced in the bloodstream of the subject in an amount effective to treat the at least one condition.

10. The method of claim 1, wherein the polypeptide is interleukin-22 (IL-22).

11. The method of claim 1, wherein the polypeptide is a cytokine.

* * * * *