

US 20240148683A1

(19) **United States**

(12) **Patent Application Publication**  
**Manku et al.**

(10) **Pub. No.: US 2024/0148683 A1**

(43) **Pub. Date:**  
**May 9, 2024**

(54) **STABLE PHARMACEUTICAL  
COMPOSITION AND METHODS OF USING  
SAME**

(71) Applicant: **Amarin Pharmaceuticals Ireland  
Limited**, Dublin (IE)

(72) Inventors: **Mehar Manku**, Birmingham (GB); **Ian  
Osterloh**, Kent (GB); **Pierre Wicker**,  
Mystic, CT (US); **Rene Braeckman**,  
Richboro, PA (US); **Paresh Soni**,  
Mystic, CT (US)

(21) Appl. No.: **18/447,880**

(22) Filed: **Aug. 10, 2023**

**Related U.S. Application Data**

(63) Continuation of application No. 17/219,039, filed on Mar. 31, 2021, which is a continuation of application No. 17/137,713, filed on Dec. 30, 2020, now Pat. No. 11,213,504, which is a continuation of application No. 16/894,567, filed on Jun. 5, 2020, now Pat. No. 10,881,632, which is a continuation of application No. 16/696,627, filed on Nov. 26, 2019, now Pat. No. 10,842,766, which is a continuation of application No. 16/562,294, filed on Sep. 5, 2019, now Pat. No. 11,147,787, which is a continuation of application No. 15/996,901, filed on Jun. 4, 2018, now Pat. No. 10,449,172, which is a continuation of application No. 15/415,468, filed on Jan. 25, 2017, now Pat. No. 10,010,517, which is a continuation of application No. 15/092,391, filed on Apr. 6, 2016, now Pat. No. 9,585,856, which is a continuation of application No. 14/709,937, filed on May 12, 2015, now abandoned, which is a continuation of application No. 14/259,724, filed on Apr. 23, 2014, now Pat. No. 9,072,715, which is a continuation of application No. 14/084,

887, filed on Nov. 20, 2013, now Pat. No. 9,060,982, which is a continuation of application No. 13/685,291, filed on Nov. 26, 2012, now Pat. No. 8,613,945, which is a continuation of application No. 13/614,111, filed on Sep. 13, 2012, now Pat. No. 8,454,994, which is a continuation of application No. 13/458,496, filed on Apr. 27, 2012, now Pat. No. 8,445,003, which is a continuation of application No. 12/769,885, filed on Apr. 29, 2010, now Pat. No. 8,298,554.

(60) Provisional application No. 61/173,763, filed on Apr. 29, 2009.

**Publication Classification**

(51) **Int. Cl.**  
*A61K 31/202* (2006.01)  
*A61K 9/48* (2006.01)  
*A61K 31/232* (2006.01)  
*A61K 45/06* (2006.01)  
*A61P 3/06* (2006.01)  
*A61P 9/00* (2006.01)  
*A61P 9/04* (2006.01)  
*A61P 9/06* (2006.01)  
*A61P 9/10* (2006.01)  
(52) **U.S. Cl.**  
CPC ..... *A61K 31/202* (2013.01); *A61K 9/48* (2013.01); *A61K 9/4825* (2013.01); *A61K 9/4858* (2013.01); *A61K 31/232* (2013.01); *A61K 45/06* (2013.01); *A61P 3/06* (2018.01); *A61P 9/00* (2018.01); *A61P 9/04* (2018.01); *A61P 9/06* (2018.01); *A61P 9/10* (2018.01)

(57) **ABSTRACT**

The present invention relates to, inter alia, pharmaceutical compositions comprising a polyunsaturated fatty acid and to methods of using the same to treat or prevent cardiovascular-related diseases.




FIG. 1




FIG. 2




FIG. 3



# STABLE PHARMACEUTICAL COMPOSITION AND METHODS OF USING SAME

## CROSS-REFERENCE TO RELATED APPLICATIONS

**[0001]** This application is a continuation of U.S. application Ser. No. 17/219,039 filed Mar. 31, 2021, which is a continuation of U.S. application Ser. No. 17/137,713 filed Dec. 30, 2020, which is a continuation of U.S. application Ser. No. 16/894,567 filed Jun. 5, 2020 (now U.S. Pat. No. 10,881,632), which is a continuation of U.S. application Ser. No. 16/696,627 filed Nov. 26, 2019 (now U.S. Pat. No. 10,842,766), which is a continuation of U.S. Application No. 16,562,294 filed Sep. 5, 2019, which is a continuation of U.S. application Ser. No. 15/996,901 filed Jun. 4, 2018 (now U.S. Pat. No. 10,449,172), which is a continuation of U.S. application Ser. No. 15/415,468 filed Jan. 25, 2017 (now U.S. Pat. No. 10,010,517), which is a continuation of U.S. application Ser. No. 15/092,391 filed Apr. 6, 2016 (now U.S. Pat. No. 9,585,856), which is a continuation of U.S. Application Ser. No. 14/709,937 filed May 12, 2015, which is a continuation of U.S. application Ser. No. 14/259,724 filed Apr. 23, 2014 (now U.S. Pat. No. 9,072,715), which is a continuation of U.S. application Ser. No. 14/084,887 filed Nov. 20, 2013 (now U.S. Pat. No. 9,060,982), which is a continuation of U.S. application Ser. No. 13/685,291 filed Nov. 26, 2012 (now U.S. Pat. No. 8,613,945), which is a continuation of U.S. application Ser. No. 13/614,111 filed Sep. 13, 2012 (now U.S. Pat. No. 8,454,994), which is a continuation of U.S. application Ser. No. 13/458,496 filed Apr. 27, 2012 (now U.S. Pat. No. 8,445,003), which is a continuation of U.S. Application Ser. No. 12/769,885 filed Apr. 29, 2010 (now U.S. Pat. No. 8,298,554), which claims priority to U.S. Provisional Application No. 61/173,763, filed Apr. 29, 2009, the entireties of each of which are incorporated herein by reference and relied upon.

## BACKGROUND

**[0002]** Mixed omega-3 fatty acid esters are typically encapsulated in type 2a gelatin capsules containing gelatin (~43.4%), glycerol (~20%) and water (~36.6%) and do not experience stability problems throughout their shelf life. While chemically modified gelatins such as succinated/succinylated gelatin have been used to encapsulate reactive fill ingredients, such gelatin is not approved for use in the U.S. and other markets.

## SUMMARY

**[0003]** We have unexpectedly found that high purity eicosapentaenoic acid (EPA) is more susceptible to oxidative degradation than mixed omega-3-acid ethyl esters. In various embodiments, the invention provides pharmaceutical compositions comprising a fatty acid or a derivative thereof in a capsule shell that resists, hinders, attenuates, or prevents oxidation of the fatty acid or fatty acid derivative, for example to a greater extent than is provided by a standard type IIa capsule shell. In a related embodiment, the fatty acid comprises eicosapentaenoic acid (EPA) or a derivative of EPA, for example ethyl eicosapentaenoate (ethyl-EPA or E-EPA). In another embodiment, the fatty acid comprises ultra-pure EPA.

**[0004]** In one embodiment, the invention provides a pharmaceutical composition comprising ultra-pure EPA encapsulated in a capsule shell, where the ultra-pure EPA has a baseline peroxide value not greater than about 5 meq/mg and upon storage of the composition at 23° C. and 50% RH for a period of time, that ultra-pure EPA has a second peroxide value not greater than about 20 meq/mg.

**[0005]** In other embodiments, the invention provides a pharmaceutical composition comprising EPA (e.g. E-EPA or ultra-pure E-EPA) encapsulated in a capsule shell comprising a film forming material and a hygroscopic plasticizer, wherein the weight ratio of film-forming material to hygroscopic plasticizer is not less than about 2.5:1. Further, the capsule shell can optionally comprise a non-hygroscopic plasticizer. In one embodiment, the capsule contains no chemically modified gelatin, for example succinated or succinylated gelatin.

**[0006]** In still other embodiments, the present invention provides methods of treating or preventing a cardiovascular-related disease using compositions as described herein.

**[0007]** These and other embodiments of the present invention will be disclosed in further detail herein below.

## BRIEF DESCRIPTION OF THE DRAWINGS

**[0008]** FIG. 1 shows dissolution profile of an inventive capsule composition containing ~500 mg E-EPA versus a composition comprising EPA in a succinated gelatin capsule.

**[0009]** FIG. 2 shows bioavailability of 300 mg of EPA in succinated gelatin capsules.

**[0010]** FIG. 3 shows bioavailability of an inventive AMR101 capsule composition containing ~500 mg E-EPA.

## DETAILED DESCRIPTION

**[0011]** While the present invention is capable of being embodied in various forms, the description below of several embodiments is made with the understanding that the present disclosure is to be considered as an exemplification of the invention, and is not intended to limit the invention to the specific embodiments illustrated. Headings are provided for convenience only and are not to be construed to limit the invention in any manner. Embodiments illustrated under any heading may be combined with embodiments illustrated under any other heading.

**[0012]** The use of numerical values in the various quantitative values specified in this application, unless expressly indicated otherwise, are stated as approximations as though the minimum and maximum values within the stated ranges were both preceded by the word "about." In this manner, slight variations from a stated value can be used to achieve substantially the same results as the stated value. Also, the disclosure of ranges is intended as a continuous range including every value between the minimum and maximum values recited as well as any ranges that can be formed by such values. Also disclosed herein are any and all ratios (and ranges of any such ratios) that can be formed by dividing a recited numeric value into any other recited numeric value. Accordingly, the skilled person will appreciate that many such ratios, ranges, and ranges of ratios can be unambiguously derived from the numerical values presented herein and in all instances such ratios, ranges, and ranges of ratios represent various embodiments of the present invention.



## Polyunsaturated Fatty Acids

**[0013]** In one embodiment, compositions of the invention comprise a polyunsaturated fatty acid as an active ingredient. In another embodiment, compositions of the invention comprise EPA as an active ingredient. The term “EPA” as used herein refers to eicosapentaenoic acid (e.g. eicosa-5,8,11,14,17-pentaenoic acid) and/or a pharmaceutically acceptable ester, derivative, conjugate or salt thereof, or mixtures of any of the foregoing.

**[0014]** In one embodiment, the EPA comprises all-cis eicosa-5,8,11,14,17-pentaenoic acid. In another embodiment, the EPA is in the form of an eicosapentaenoic acid ester. In another embodiment, the EPA comprises a  $C_{1-C_5}$  alkyl ester of EPA. In another embodiment, the EPA comprises eicosapentaenoic acid ethyl ester, eicosapentaenoic acid methyl ester, eicosapentaenoic acid propyl ester, or eicosapentaenoic acid butyl ester. In still another embodiment, the EPA comprises all-cis eicosa-5,8,11,14,17-pentaenoic acid ethyl ester.

**[0015]** In still other embodiments, the EPA comprises ethyl-EPA, lithium EPA, mono, di- or triglyceride EPA or any other ester or salt of EPA, or the free acid form of EPA. The EPA may also be in the form of a 2-substituted derivative or other derivative which slows down its rate of oxidation but does not otherwise change its biological action to any substantial degree.

**[0016]** The term “pharmaceutically acceptable” in the present context means that the substance in question does not produce unacceptable toxicity to the subject or interaction with other components of the composition.

**[0017]** In one embodiment, EPA present in a composition of the invention comprises ultra-pure EPA. The term “ultra-pure” as used herein with respect to EPA refers to a composition comprising at least 96% by weight EPA (as the term “EPA” is defined and exemplified herein). Ultra-pure EPA can comprise even higher purity EPA, for example at least 97% by weight EPA or at least 98% by weight EPA, wherein the EPA is any form of EPA as set forth herein. Ultra-pure EPA can further be defined (e.g. impurity profile) by any of the description of EPA provided herein.

**[0018]** In other embodiments, EPA is present in a composition of the invention in an amount of about 50 mg to about 5000 mg, about 75 mg to about 2500 mg, or about 100 mg to about 1000 mg, for example about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, about 300 mg, about 325 mg, about 350 mg, about 375 mg, about 400 mg, about 425 mg, about 450 mg, about 475 mg, about 500 mg, about 525 mg, about 550 mg, about 575 mg, about 600 mg, about 625 mg, about 650 mg, about 675 mg, about 700 mg, about 725 mg, about 750 mg, about 775 mg, about 800 mg, about 825 mg, about 850 mg, about 875 mg, about 900 mg, about 925 mg, about 950 mg, about 975 mg, about 1000 mg, about 1025 mg, about 1050 mg, about 1075 mg, about 1100 mg, about 1025 mg, about 1050 mg, about 1075 mg, about 1200 mg, about 1225 mg, about 1250 mg, about 1275 mg, about 1300 mg, about 1325 mg, about 1350 mg, about 1375 mg, about 1400 mg, about 1425 mg, about 1450 mg, about 1475 mg, about 1500 mg, about 1525 mg, about 1550 mg, about 1575 mg, about 1600 mg, about 1625 mg, about 1650 mg, about 1675 mg, about 1700 mg, about 1725 mg, about 1750 mg, about 1775 mg, about 1800 mg, about 1825 mg, about 1850 mg, about 1875 mg, about 1900 mg, about 1925 mg, about 1950 mg, about 1975 mg, about 2000 mg, about

2025 mg, about 2050 mg, about 2075 mg, about 2100 mg, about 2125 mg, about 2150 mg, about 2175 mg, about 2200 mg, about 2225 mg, about 2250 mg, about 2275 mg, about 2300 mg, about 2325 mg, about 2350 mg, about 2375 mg, about 2400 mg, about 2425 mg, about 2450 mg, about 2475 mg, or about 2500 mg.

**[0019]** In various embodiments, one or more antioxidants can be present in the EPA (e.g. E-EPA or ultra-pure E-EPA). Non-limiting examples of suitable antioxidants include tocopherol, lecithin, citric acid and/or ascorbic acid. One or more antioxidants, if desired, are typically present in the EPA in an amount of about 0.01% to about 0.1%, by weight, or about 0.025% to about 0.05%, by weight.

**[0020]** In one embodiment, a composition of the invention contains not more than about 10%, not more than about 9%, not more than about 8%, not more than about 7%, not more than about 6%, not more than about 5%, not more than about 4%, not more than about 3%, not more than about 2%, not more than about 1%, or not more than about 0.5%, by weight of total fatty acids, docosahexaenoic acid or derivative thereof such as E-DHA, if any. In another embodiment, a composition of the invention contains substantially no docosahexaenoic acid or derivative thereof such as E-DHA. In still another embodiment, a composition of the invention contains no docosahexaenoic acid or E-DHA.

**[0021]** In another embodiment, EPA represents at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, at least about 98%, at least about 99%, or 100%, by weight, of all fatty acids present in a composition of the invention.

**[0022]** In another embodiment, a composition of the invention contains less than 30%, less than 20%, less than 10%, less than 9%, less than 8%, less than 7%, less than 6%, less than 5%, less than 4%, less than 3%, less than 2%, less than 1%, less than 0.5% or less than 0.25%, by weight of the total composition or by weight of the total fatty acid content, of any fatty acid other than EPA, or derivative thereof. Illustrative examples of a “fatty acid other than EPA” include linolenic acid (LA) or derivative thereof such as ethyl-linolenic acid, arachidonic acid (AA) or derivative thereof such as ethyl-AA, docosahexaenoic acid (DHA) or derivative thereof such as ethyl-DHA, alpha-linolenic acid (ALA) or derivative thereof such as ethyl-ALA, stearadonic acid (STA) or derivative thereof such as ethyl-SA, eicosatrienoic acid (ETA) or derivative thereof such as ethyl-ETA and/or docosapentaenoic acid (DPA) or derivative thereof such as ethyl-DPA.

**[0023]** In another embodiment, a composition of the invention has one or more of the following features: (a) eicosapentaenoic acid ethyl ester represents at least 96%, at least 97%, or at least 98%, by weight, of all fatty acids present in the composition; (b) the composition contains not more than 4%, not more than 3%, or not more than 2%, by weight, of total fatty acids other than eicosapentaenoic acid ethyl ester; (c) the composition contains not more than 0.6%, 0.5%, or 0.4% of any individual fatty acid other than eicosapentaenoic acid ethyl ester; (d) the composition has a refractive index (20° C.) of about 1 to about 2, about 1.2 to about 1.8 or about 1.4 to about 1.5; (e) the composition has a specific gravity (20° C.) of about 0.8 to about 1.0, about 0.85 to about 0.95 or about 0.9 to about 0.92; (f) the composition contains not more than 20 ppm, 15 ppm or 10 ppm heavy metals, (g) the composition contains not more



than 5 ppm, 4 ppm, 3 ppm, or 2 ppm arsenic, and/or (h) the composition has a peroxide value not more than 5, 4, 3, or 2.

**[0024]** In another embodiment, a composition useful in accordance with the invention comprises, consists essentially of or consists of at least 95% by weight ethyl eicosapentaenoate (EPA-E), about 0.2% to about 0.5% by weight ethyl octadecatetraenoate (ODTA-E), about 0.05% to about 0.25% by weight ethyl nonaecapentaenoate (NDPA-E), about 0.2% to about 0.45% by weight ethyl arachidonate (AA-E), about 0.3% to about 0.5% by weight ethyl eicosatetraenoate (ETA-E), and about 0.05% to about 0.32% ethyl heneicosapentaenoate (HPA-E). In another embodiment, the composition is present in a capsule shell. In still another embodiment, the capsule shell contains no chemically modified gelatin.

**[0025]** In another embodiment, compositions useful in accordance with the invention comprise, consist essentially of, or consist of at least 95%, 96% or 97%, by weight, ethyl eicosapentaenoate, about 0.2% to about 0.5% by weight ethyl octadecatetraenoate, about 0.05% to about 0.25% by weight ethyl nonaecapentaenoate, about 0.2% to about 0.45% by weight ethyl arachidonate, about 0.3% to about 0.5% by weight ethyl eicosatetraenoate, and about 0.05% to about 0.32% by weight ethyl heneicosapentaenoate. Optionally, the composition contains not more than about 0.06%, about 0.05%, or about 0.04%, by weight, DHA or derivative thereof such as ethyl-DHA. In one embodiment the composition contains substantially no or no amount of DHA or derivative thereof such as ethyl-DHA. The composition further optionally comprises one or more antioxidants (e.g. tocopherol) in an amount of not more than about 0.5% or not more than 0.05%. In another embodiment, the composition comprises about 0.05% to about 0.4%, for example about 0.2% by weight tocopherol. In another embodiment, about 500 mg to about 1 g of the composition is provided in a capsule shell. In another embodiment, the capsule shell contains no chemically modified gelatin.

**[0026]** In another embodiment, compositions useful in accordance with the invention comprise, consist essentially of, or consist of at least 96% by weight ethyl eicosapentaenoate, about 0.22% to about 0.4% by weight ethyl octadecatetraenoate, about 0.075% to about 0.20% by weight ethyl nonaecapentaenoate, about 0.25% to about 0.40% by weight ethyl arachidonate, about 0.3% to about 0.4% by weight ethyl eicosatetraenoate and about 0.075% to about 0.25% by weight ethyl heneicosapentaenoate. Optionally, the composition contains not more than about 0.06%, about 0.05%, or about 0.04%, by weight, DHA or derivative thereof such as ethyl-DHA. In one embodiment the composition contains substantially no or no amount of DHA or derivative thereof such as ethyl-DHA. The composition further optionally comprises one or more antioxidants (e.g. tocopherol) in an amount of not more than about 0.5% or not more than 0.05%. In another embodiment, the composition comprises about 0.05% to about 0.4%, for example about 0.2% by weight tocopherol. In another embodiment, the invention provides a dosage form comprising about 500 mg to about 1 g of the foregoing composition in a capsule shell. In one embodiment, the dosage form is a gel- or liquid-containing capsule and is packaged in blister packages of about 1 to about 20 capsules per sheet.

**[0027]** In another embodiment, compositions useful in accordance with the invention comprise, consist essentially

of or consist of at least 96%, 97% or 98%, by weight, ethyl eicosapentaenoate, about 0.25% to about 0.38% by weight ethyl octadecatetraenoate, about 0.10% to about 0.15% by weight ethyl nonaecapentaenoate, about 0.25% to about 0.35% by weight ethyl arachidonate, about 0.31% to about 0.38% by weight ethyl eicosatetraenoate, and about 0.08% to about 0.20% by weight ethyl heneicosapentaenoate. Optionally, the composition contains not more than about 0.06%, about 0.05%, or about 0.04%, by weight, DHA or derivative thereof such as ethyl-DHA. In one embodiment the composition contains substantially no or no amount of DHA or derivative thereof such as ethyl-DHA. The composition further optionally comprises one or more antioxidants (e.g. tocopherol) in an amount of not more than about 0.5% or not more than 0.05%. In another embodiment, the composition comprises about 0.05% to about 0.4%, for example about 0.2% by weight tocopherol. In another embodiment, the invention provides a dosage form comprising about 500 mg to about 1 g of the foregoing composition in a capsule shell. In another embodiment, the capsule shell contains no chemically modified gelatin.

**[0028]** In various embodiments, the invention provides a polyunsaturated fatty acid such as EPA (e.g. E-EPA or ultra-pure E-EPA) encapsulated in a pharmaceutical capsule shell. In one embodiment, the capsule shell resists, hinders, attenuates, or prevents oxidation of the fatty acid or fatty acid derivative. In another embodiment, the capsule shell resists, hinders, attenuates, or prevents oxidation of the polyunsaturated fatty acid or derivative to a greater extent than a standard type IIa gelatin capsule. In another embodiment, the capsule contains no chemically modified gelatin, for example succinated, succinylated, phthalated, carbamylated and/or phenol carbamylated gelatin.

**[0029]** In another embodiment, the invention provides a pharmaceutical composition comprising EPA (e.g. E-EPA or ultra-pure E-EPA) encapsulated in a capsule shell as described herein and having a baseline peroxide value not greater than about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg, wherein upon storage of the composition at 23° C. and 50% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, the ultra-pure EPA has a second peroxide value not greater than about 25 meq/mg, about 24 meq/mg, about 23 meq/mg, about 22 meq/mg, about 21 meq/mg, about 20 meq/mg, about 19 meq/mg, about 18 meq/mg, about 17 meq/mg, about 16 meq/mg, about 15 meq/mg, about 14 meq/mg, about 13 meq/mg, about 12 meq/mg, about 11 meq/mg, about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg.

**[0030]** The “baseline peroxide value” and “second peroxide values” can be measured in any suitable manner, for example by using a U.S. or PhEur or JP compendial method. Typically, a plurality of encapsulated EPA compositions are provided, each composition containing EPA having been encapsulated at substantially the same time. A first sampling of 1 or more capsules from the plurality is provided, the capsules are opened and peroxide value of the EPA is measured substantially immediately thereafter, providing an average baseline peroxide value. At substantially the same



time, a second sampling of 1 or more capsules from the plurality are provided and are placed under desired storage conditions for a desired time period. At the end of the desired time period, the capsules are opened and peroxide value of the EPA is measured substantially immediately thereafter, providing an average second peroxide value. The baseline and second peroxide values can then be compared. In one embodiment, the “baseline peroxide value” and “second peroxide value” are determined using a plurality of encapsulated EPA dosage units wherein each dosage unit was encapsulated (i.e. the EPA filled and sealed into capsules) within a same 60 day period, same 30 day period, a same 20 day period, a same 10 day period, a same 5 day period or a same 1 day period.

**[0031]** In another embodiment, the invention provides a pharmaceutical composition comprising EPA (e.g. E-EPA or ultra-pure E-EPA) encapsulated in a capsule shell as described herein and having a baseline peroxide value not greater than about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg, wherein upon storage of the composition at 25° C. and 60% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition has a second peroxide value not greater than about 25 meq/mg, about 24 meq/mg, about 23 meq/mg, about 22 meq/mg, about 21 meq/mg, about 20 meq/mg, about 19 meq/mg, about 18 meq/mg, about 17 meq/mg, about 16 meq/mg, about 15 meq/mg, about 14 meq/mg, about 13 meq/mg, about 12 meq/mg, about 11 meq/mg, about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg.

**[0032]** In another embodiment, the invention provides a pharmaceutical composition comprising EPA (e.g. E-EPA or ultra-pure E-EPA) encapsulated in a capsule shell as described herein and having a baseline peroxide value not greater than about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg, wherein upon storage of the composition at 30° C. and 65% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition has a second peroxide value not greater than about 25 meq/mg, about 24 meq/mg, about 23 meq/mg, about 22 meq/mg, about 21 meq/mg, about 20 meq/mg, about 19 meq/mg, about 18 meq/mg, about 17 meq/mg, about 16 meq/mg, about 15 meq/mg, about 14 meq/mg, about 13 meq/mg, about 12 meq/mg, about 11 meq/mg, about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg.

**[0033]** In another embodiment, the invention provides a pharmaceutical composition comprising EPA (e.g. E-EPA or ultra-pure E-EPA) encapsulated in a capsule shell as described herein and having a baseline peroxide value not greater than about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg, wherein upon storage of the composition at 40° C. and 75%

RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition has a second peroxide value not greater than about 25 meq/mg, about 24 meq/mg, about 23 meq/mg, about 22 meq/mg, about 21 meq/mg, about 20 meq/mg, about 19 meq/mg, about 18 meq/mg, about 17 meq/mg, about 16 meq/mg, about 15 meq/mg, about 14 meq/mg, about 13 meq/mg, about 12 meq/mg, about 11 meq/mg, about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg.

**[0034]** In another embodiment, the invention provides a pharmaceutical composition comprising EPA (e.g. E-EPA or ultra-pure E-EPA) encapsulated in a capsule shell and having a baseline peroxide value not greater than about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg, wherein the capsule comprises a film-forming material and a plasticizer in a weight ratio of not less than 1.75:1 and wherein upon storage of the composition at 23° C. and 50% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition has a second peroxide value not greater than about 25 meq/mg, about 24 meq/mg, about 23 meq/mg, about 22 meq/mg, about 21 meq/mg, about 20 meq/mg, about 19 meq/mg, about 18 meq/mg, about 17 meq/mg, about 16 meq/mg, about 15 meq/mg, about 14 meq/mg, about 13 meq/mg, about 12 meq/mg, about 11 meq/mg, about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg.

**[0035]** In another embodiment, the invention provides a pharmaceutical composition comprising EPA (e.g. E-EPA or ultra-pure E-EPA) encapsulated in a capsule shell and having a baseline peroxide value not greater than about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg, wherein the capsule comprises a film-forming material and a plasticizer in a weight ratio of not less than 1.75:1 and wherein upon storage of the composition at 25° C. and 60% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition has a second peroxide value not greater than about 25 meq/mg, about 24 meq/mg, about 23 meq/mg, about 22 meq/mg, about 21 meq/mg, about 20 meq/mg, about 19 meq/mg, about 18 meq/mg, about 17 meq/mg, about 16 meq/mg, about 15 meq/mg, about 14 meq/mg, about 13 meq/mg, about 12 meq/mg, about 11 meq/mg, about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg.

**[0036]** In another embodiment, the invention provides a pharmaceutical composition comprising EPA (e.g. E-EPA or ultra-pure E-EPA) encapsulated in a capsule shell and having a baseline peroxide value not greater than about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg,



mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg, wherein the capsule comprises a film-forming material and a plasticizer in a weight ratio of not less than 1.75:1 and wherein upon storage of the composition at 30° C. and 65% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition has a second peroxide value not greater than about 25 meq/mg, about 24 meq/mg, about 23 meq/mg, about 22 meq/mg, about 21 meq/mg, about 20 meq/mg, about 19 meq/mg, about 18 meq/mg, about 17 meq/mg, about 16 meq/mg, about 15 meq/mg, about 14 meq/mg, about 13 meq/mg, about 12 meq/mg, about 11 meq/mg, about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg.

**[0037]** In another embodiment, the invention provides a pharmaceutical composition comprising EPA (e.g. E-EPA or ultra-pure E-EPA) encapsulated in a capsule shell and having a baseline peroxide value not greater than about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg, wherein the capsule comprises a film-forming material and a plasticizer in a weight ratio of not less than 1.75:1 and wherein upon storage of the composition at 40° C. and 75% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition has a second peroxide value not greater than about 25 meq/mg, about 24 meq/mg, about 23 meq/mg, about 22 meq/mg, about 21 meq/mg, about 20 meq/mg, about 19 meq/mg, about 18 meq/mg, about 17 meq/mg, about 16 meq/mg, about 15 meq/mg, about 14 meq/mg, about 13 meq/mg, about 12 meq/mg, about 11 meq/mg, about 10 meq/mg, about 9 meq/mg, about 8 meq/mg, about 7 meq/mg, about 6 meq/mg, about 5 meq/mg, about 4 meq/mg, about 3 meq/mg or about 2 meq/mg.

**[0038]** In another embodiment, the invention provides a pharmaceutical composition comprising encapsulated EPA (e.g. E-EPA or ultra-pure E-EPA) containing a labeled amount (i.e. initial amount) of EPA or E-EPA, wherein upon storage of the composition at 23° C. and 50% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition contains at least about 97%, about 98%, about 99%, about 99.5%, about 99.7%, about 99.9% or substantially all or 100% of the labeled amount of EPA or E-EPA, by weight.

**[0039]** In another embodiment, the invention provides a pharmaceutical composition comprising encapsulated EPA (e.g. E-EPA or ultra-pure E-EPA) containing a labeled amount (i.e. initial amount) of EPA or E-EPA, wherein upon storage of the composition at 25° C. and 60% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition contains at least about 97%, about

98%, about 99%, about 99.5%, about 99.7%, about 99.9% or substantially all or 100% of the labeled amount of EPA or E-EPA, by weight.

**[0040]** In another embodiment, the invention provides a pharmaceutical composition comprising encapsulated EPA (e.g. E-EPA or ultra-pure E-EPA) containing a labeled amount of EPA or E-EPA, wherein upon storage of the composition at 30° C. and 65% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition contains at least about 97%, about 98%, about 99%, about 99.5%, about 99.7%, about 99.9%, substantially all or 100% of the labeled amount of EPA or E-EPA, by weight.

**[0041]** In another embodiment, the invention provides a pharmaceutical composition comprising encapsulated EPA (e.g. E-EPA or ultra-pure E-EPA) containing a labeled amount of EPA, wherein upon storage of the composition at 40° C. and 75% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition contains at least about 97%, about 98%, about 99%, about 99.5%, about 99.7%, about 99.8%, about 99.9%, substantially all or 100% of the labeled amount of EPA or E-EPA, by weight.

**[0042]** In another embodiment, the invention provides a pharmaceutical composition comprising encapsulated EPA containing a labeled amount of EPA or E-EPA, wherein upon storage of the composition at 23° C. and 50% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition contains not more than about 0.5%, not more than about 0.25%, not more than about 0.15%, not more than about 0.125%, not more than about 0.1%, not more than about 0.075%, not more than about 0.05% or substantially no degradation product and/or specified degradation product. The term “degradation product” in the present context means “an impurity resulting from a chemical change in the composition brought about during manufacture and/or storage of the composition by the effect of, for example, light, temperature, pH, water or by reaction with an excipient and/or the immediate container closure system.” The term “specified degradation product in the present context means “a degradation product, either identified or unidentified, that is individually listed and limited with a specific acceptance criterion in the product specification” for a particular product.

**[0043]** In another embodiment, the invention provides a pharmaceutical composition comprising encapsulated EPA (e.g. E-EPA or ultra-pure E-EPA) containing a labeled amount of EPA or E-EPA, wherein upon storage of the composition at 25° C. and 60% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, the composition contains not more than about 0.5%, not more than about 0.25%, not more than about 0.15%, not more than about 0.125%, not more than about 0.1%, not more than







22, about 23 or about 24 months, said composition contains not more than about 0.5% (by weight of the labeled EPA or E-EPA), not more than about 0.25%, not more than about 0.15%, not more than about 0.125%, not more than about 0.1%, not more than about 0.075%, not more than about 0.05% or substantially no degradation product and/or specified degradation product.

**[0052]** In another embodiment, the invention provides a pharmaceutical composition comprising encapsulated (e.g. E-EPA or ultra-pure E-EPA) containing a labeled amount of EPA or E-EPA, wherein the capsule comprises a film-forming material and a plasticizer in a weight ratio of about 2:5:1 to about 10:1 and upon storage of the composition at 30° C. and 65% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition contains not more than about 0.5% (by weight of the labeled EPA or E-EPA), not more than about 0.25%, not more than about 0.15%, not more than about 0.125%, not more than about 0.1%, not more than about 0.075%, not more than about 0.05% or substantially no degradation product and/or specified degradation product.

**[0053]** In another embodiment, the invention provides a pharmaceutical composition comprising encapsulated (e.g. E-EPA or ultra-pure E-EPA) containing a labeled amount of EPA or E-EPA, wherein the capsule comprises a film-forming material and a plasticizer in a weight ratio of about 2:5:1 to about 10:1 and upon storage of the composition at 40° C. and 75% RH for a period about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 months, said composition contains not more than about 0.5% (by weight of the labeled EPA or E-EPA), not more than about 0.25%, not more than about 0.15%, not more than about 0.125%, not more than about 0.1%, not more than about 0.075%, not more than about 0.05% or substantially no degradation product and/or specified degradation product.

**[0054]** In another embodiment, the present invention provides a pharmaceutical composition comprising about 0.5 g to about 1.5 g of EPA (e.g. E-EPA or ultra-pure E-EPA) having a labeled amount of EPA or E-EPA encapsulated in a pharmaceutical capsule, wherein upon storage at 15° C. to 30° C. for a period of about 6 months, 12 months, 18 months, 24 months, 30 months, or 36 months, at least about 97%, about 98%, about 99%, about 99.5%, about 99.6%, about 99.7%, about 99.8%, about 99.9% or substantially all of the labeled amount of EPA is still present in the composition. In a related embodiment, the composition has not reached its labeled expiration date during said storage period.

**[0055]** In various embodiments, capsule shells suitable for use in the present invention comprise one or more film-forming materials, one or more plasticizers and optionally a solvent (e.g. water). In a related embodiment, the film-forming material comprises gelatin. In another embodiment, the plasticizer comprises a hygroscopic and/or non-hygroscopic plasticizer. In still another embodiment, the capsule shell comprises a film-forming material, a hygroscopic plasticizer, a non-hygroscopic plasticizer and a solvent.

**[0056]** In another embodiment, the capsule shell comprises about 30% to about 70% or about 40% to about 65%, by weight, of a film-forming material, about 15% to about 40% or about 20% to about 35%, by weight, of one or more plasticizers, and about 3% to about 15% or about 5% to about 10%, by weight, solvent such as water. Optionally, the capsules may also contain additives such as colorants, flavorants, preservatives, disintegrants, surfactants, fragrances, sweeteners, etc.

**[0057]** Capsules suitable for use in various embodiments of the invention comprise a film-forming material, for example gelatin. Gelatin is typically manufactured from animal byproducts that contain collagen, for example in the bones, skin, and connective tissue. Methods of producing gelatin from animal byproducts are well-known in the art. In various embodiments, the gelatin may be alkali-treated gelatin, acid-treated gelatin, chemically modified gelatin, or mixtures thereof. Methods to produce alkali-treated gelatin, acid-treated gelatin, and chemically modified gelatin are known in the art and are described, for example in Nakamura et al., U.S. 2003/0195246, hereby incorporated by reference herein in its entirety.

**[0058]** The film-forming material may also comprise, for example, non-animal based hydrocolloids such as carrageenan, alkylated or hydroxyalkylated cellulose ethers, starch, alpha-starch, hydroxyalkyl starch, sodium alginate, sodium salt of a gelatin copolymer and acrylic acid.

**[0059]** In another embodiment, the film-forming material can comprise a 20:80 to about 80:20, by weight, mixture, for example a 60:40, by weight mixture of hydroxypropyl methyl cellulose and polyvinyl alcohol (e.g. about 70% to about 90%, for example about 88.0% saponified; and about 30 to about 50, for example about 45.0 centipoise viscosity). In another embodiment, the film-forming material can comprise a 20:80 to about 80:20, by weight, mixture, for example a 60:40, by weight, mixture of hydroxyethyl cellulose and polyvinyl alcohol (e.g. about 70% to about 99.9%, for example about 98.5% saponified; and about 2 to about 30, for example about 5.5 centipoise viscosity).

**[0060]** A suitable capsule shell may further comprise an elasticity reducing gel extender as part of the film-forming material. An elasticity reducing gel extender can comprise starch, starch derivatives such as high amylose starch, oxidized starch, esterified starch, acid-thinned starch, etherified starch, hydrolyzed starch, hydrolyzed and hydrogenated starch, enzyme treated starch, and modified celluloses or other natural or modified natural biopolymers such as bacterial polysaccharides, vegetable gums, or other exudates including alginates, carrageenans, guar gum, gum arabic, gum ghatti, gum karaya, gum tragacanth, pectins, tamarind gum, xanthan gum, and dextrans as well as synthetic polymers such as carbon chain polymers of the vinyl and acrylic types as well as heterochains of the polyoxide and polyamine types including polyethylene oxide, polypropylene oxide, polyoxymethylene, polytrimethylene oxide, block copolymers of ethylene oxide, block copolymers of polyethylene oxide, polyvinyl methyl ether, polyethylene imine, polyacrylic acid, polyacrylamide, polymethacrylic acid, polymethacrylamide, poly(N,N-Dimethylacrylamide), poly(N-Isopropylacrylamide), poly(N-Acrylylglycinamide), poly(N-Methacrylyglycinamide), acrylic copolymers, polyvinyl alcohol polyvinylacetate, polyvinyl acetate-co-vinyl alcohol, polyvinylpyrrolidone, N-Methylpyrrolidone, N-Ethylpyrrolidone, N-Vinylpyrrolidone, sarcosine anhy-



dride, polyvinylloxazolidone, and polyvinylmethyloxazolidone. The starch or other elasticity reducing gel extender may be added into the formulation in amounts ranging from about 8% to about 30% by weight, for example about 10% to about 16%, by weight.

**[0061]** Capsule shells suitable for use in various embodiments of the invention can comprise one or more plasticizers, for example hygroscopic and/or non-hygroscopic plasticizers. Non-limiting examples of suitable hygroscopic plasticizers include glycerin, sorbitol and alkylene glycols (e.g., propylene glycol and low molecular weight polyethylene glycols). Non-limiting examples of suitable non-hygroscopic plasticizers include partially dehydrated hydrogenated glucose syrup, maltitol, maltose, lactitol, xylitol, erythritol and polyethylene glycols of average molecular weights from about 400 to about 6000.

**[0062]** In one embodiment, a capsule shell suitable for use in a composition of the invention has a hygroscopic plasticizer to non-hygroscopic plasticizer weight ratio of about 1:1 to about 8:1, about 2:1 to about 6:1, about 3:1 to about 5:1, for example about 4:1, about 4.25:1, about 4.5:1 or about 4.75:1.

**[0063]** In another embodiment, a capsule shell suitable for use in a composition of the invention has a gelatin to glycerol weight ratio of about 2:5:1 to about 10:1, about 3.5:1 to about 9:1, about 4:1 to about 8:1, or about 5:1 to about 7:1, for example at least about 2.6:1, at least about 2.7:1, at least about 2.8:1, at least about 2.9:1, at least about 3:1, at least about 3.1:1, at least about 3.2:1, at least about 3.3:1, at least about 3.4:1, at least about 3.5:1, at least about 3.6:1, at least about 3.7:1, at least about 3.8:1, at least about 3.9:1, at least about 4.0:1, at least about 4.1:1, at least about 4.2:1, at least about 4.3:1, at least about 4.4:1, at least about 4.5:1, at least about 4.6:1, at least about 4.7:1, at least about 4.8:1, at least about 4.9:1, at least about 5.0:1, at least about 5.1:1, or at least about 5.2:1.

**[0064]** In another embodiment, a suitable capsule shell has a film-forming material (e.g. gelatin) to total plasticizer weight ratio of about 1.75 to about 5, about 1.78 to about 3, or about 1.8 to about 2.5, for example at least about 1.76, at least about 1.77, at least about 1.78, at least about 1.79, at least about 1.8, at least about 1.81, at least about 1.82, at least about 1.83, or at least about 1.84.

**[0065]** In another embodiment, the capsule shell has: (1) a gelatin to glycerol weight ratio of about 2:5:1 to about 10:1, about 3.5:1 to about 9:1, about 4:1 to about 8:1, or about 5:1 to about 7:1, for example at least about 2.6:1, at least about 2.7:1, at least about 2.8:1, at least about 2.9:1, at least about 3:1, at least about 3.1:1, at least about 3.2:1, at least about 3.3:1, at least about 3.4:1, at least about 3.5:1, at least about 3.6:1, at least about 3.7:1, at least about 3.8:1, at least about 3.9:1, at least about 4.0:1, at least about 4.1:1, at least about 4.2:1, at least about 4.3:1, at least about 4.4:1, at least about 4.5:1, at least about 4.6:1, at least about 4.7:1, at least about 4.8:1, at least about 4.9:1, at least about 5.0:1, at least about 5.1:1, or at least about 5.2:1; and/or (2) a gelatin to total plasticizer weight ratio of about 1.75:1 to about 5:1, about 1.78:1 to about 3:1, or about 1.8:1 to about 2.5:1, for example at least about 1.76:1, at least about 1.77:1, at least about 1.78:1, at least about 1.79:1, at least about 1.8:1, at least about 1.81, at least about 1.82, at least about 1.83, or at least about 1.84.

**[0066]** In one embodiment, the capsule shell comprises one or more of: gelatin in an amount of about 50% to about

70%; glycerol in an amount of about 5% to about 15%; sorbitol in an amount of about 15% to about 25%; and/or maltitol in an amount of about 3% to about 10%, by weight of the non-aqueous components. Such a capsule can further comprise about 2% to about 16% by weight of a solvent such as water.

**[0067]** In another embodiment, a capsule shell suitable for use in compositions of the present invention can be prepared using a gel mass comprising about 40% to about 50% gelatin, about 2% to about 12% glycerol, about 10% to about 20% sorbitol solution, about 2% to about 10% maltitol syrup, and about 20% to about 35% water, by weight. In one embodiment, a capsule shell suitable for use in a composition of the present invention can be prepared using a gel mass comprising about 45% gelatin by weight, about 7% glycerol by weight, about 17% sorbitol solution (e.g. 30% water) by weight, about 6% maltitol syrup (e.g. 15%-32% water) by weight, and about 25% water by weight. Capsules prepared from such a gel mass can be dried to about 2% to about 12% final moisture content. Capsules prepared by such a process that contain EPA (e.g. E-EPA or ultra-pure E-EPA), and methods of using the same in the treatment of cardiovascular-related diseases represent further embodiments of the invention. Capsule compositions as described herein can further comprise coatings, for example enteric polymer or wax coatings.

**[0068]** In one embodiment, a composition of the invention provides a relatively rapid dissolution profile yet still maintains excellent stability of the encapsulated material (e.g. EPA). In a related embodiment, a composition of the invention has a dissolution profile (as measured by Rotating Dialysis Cell Dissolution (RDC) Apparatus under the conditions set forth herein below) of one or more of the following: (1) at least about 20%, at least about 23% or at least about 25% of E-EPA is dissolved by 10 minutes; (2) at least about 45%, at least about 50% or at least about 55% of E-EPA is dissolved by 30 minutes; (3) at least about 80%, at least about 82%, at least about 85% or at least about 87% of E-EPA is dissolved by 60 minutes; and/or (4) at least about 95%, at least about 97% or 100% of E-EPA is dissolved by 100 minutes. In a related embodiment, the fill material still retains the stability/peroxide values as set forth throughout this specification.

**[0069]** In another embodiment, a composition of the invention provides a relatively short  $T_{max}$  yet still maintains excellent stability of the encapsulated material (e.g. EPA). In a related embodiment, a composition of the invention, upon administration to a subject, exhibits an EPA  $T_{max}$  less than 6 hours, less than 5.8 hours, less than 5.6 hours, less than 5.4 hours or less than 5.2 hours, for example about 4.8 to about 5.2 hours. In a related embodiment, the fill material still retains the stability/peroxide values as set forth throughout this specification.

**[0070]** In one embodiment, a method for treatment and/or prevention of a cardiovascular-related disease using a composition as described herein is provided. The term “cardiovascular-related disease” herein refers to any disease or disorder of the heart or blood vessels (i.e. arteries and veins) or any symptom thereof. The term “cardiovascular-related disease” herein refers to any disease or disorder of the heart or blood vessels (i.e. arteries and veins) or any symptom thereof, or any disease or condition that causes or contributes to a cardiovascular disease.” Non-limiting examples of cardiovascular-related diseases include acute cardiac isch-



emic events, acute myocardial infarction, angina, angina pectoris, arrhythmia, atrial fibrillation, atherosclerosis, arterial fibrillation, cardiac insufficiency, cardiovascular disease, chronic heart failure, chronic stable angina, congestive heart failure, coronary artery disease, coronary heart disease, deep vein thrombosis, diabetes, diabetes mellitus, diabetic neuropathy, diastolic dysfunction in subjects with diabetes mellitus, edema, essential hypertension, eventual pulmonary embolism, fatty liver disease, heart disease, heart failure, homozygous familial hypercholesterolemia (HoFH), homozygous familial sitosterolemia, hypercholesterolemia, hyperlipidemia, hyperlipidemia in HIV positive subjects, hypertension, hypertriglyceridemia, ischemic complications in unstable angina and myocardial infarction, low blood pressure, metabolic syndrome, mixed dyslipidemia, moderate to mild heart failure, myocardial infarction, obesity management, paroxysmal atrial/arterial fibrillation/fibrillation/flutter, paroxysmal supraventricular tachycardias (PSVT), particularly severe or rapid onset edema, platelet aggregation, primary hypercholesterolemia, primary hyperlipidemia, pulmonary arterial hypertension, pulmonary hypertension, recurrent hemodynamically unstable ventricular tachycardia (VT), recurrent ventricular arrhythmias, recurrent ventricular fibrillation (VF), ruptured aneurysm, sitosterolemia, stroke, supraventricular tachycardia, symptomatic atrial fibrillation/flutter, tachycardia, type-II diabetes, vascular disease, venous thromboembolism, ventricular arrhythmias, and other cardiovascular events.

**[0071]** The term “treatment” in relation a given disease or disorder, includes, but is not limited to, inhibiting the disease or disorder, for example, arresting the development of the disease or disorder; relieving the disease or disorder, for example, causing regression of the disease or disorder; or relieving a condition caused by or resulting from the disease or disorder, for example, relieving, preventing or treating symptoms of the disease or disorder. The term “prevention” in relation to a given disease or disorder means: preventing the onset of disease development if none had occurred, preventing the disease or disorder from occurring in a subject that may be predisposed to the disorder or disease but has not yet been diagnosed as having the disorder or disease, and/or preventing further disease/disorder development if already present.

**[0072]** In one embodiment, the present invention provides a method of blood lipid therapy comprising administering to a subject or subject group in need thereof a pharmaceutical composition as described herein. In another embodiment, the subject or subject group has hypertriglyceridemia, hypercholesterolemia, mixed dyslipidemia and/or very high triglycerides.

**[0073]** In another embodiment, the subject or subject group being treated has a baseline triglyceride level (or median baseline triglyceride level in the case of a subject group), fed or fasting, of at least about 300 mg/dl, at least about 400 mg/dl, at least about 500 mg/dl, at least about 600 mg/dl, at least about 700 mg/dl, at least about 800 mg/dl, at least about 900 mg/dl, at least about 1000 mg/dl, at least about 1100 mg/dl, at least about 1200 mg/dl, at least about 1300 mg/dl, at least about 1400 mg/dl, or at least about 1500 mg/dl, for example about 400 mg/dl to about 2500 mg/dl, about 450 mg/dl to about 2000 mg/dl or about 500 mg/dl to about 1500 mg/dl.

**[0074]** In another embodiment, the subject or subject group being treated in accordance with methods of the

invention has previously been treated with Lovaza® and has experienced an increase in, or no decrease in, LDL-C levels and/or non-HDL-C levels. In one such embodiment, Lovaza® therapy is discontinued and replaced by a method of the present invention.

**[0075]** In another embodiment, the subject or subject group being treated in accordance with methods of the invention exhibits a fasting baseline absolute plasma level of free EPA (or mean thereof in the case of a subject group) not greater than about 0.70 nmol/ml, not greater than about 0.65 nmol/ml, not greater than about 0.60 nmol/ml, not greater than about 0.55 nmol/ml, not greater than about 0.50 nmol/ml, not greater than about 0.45 nmol/ml, or not greater than about 0.40 nmol/ml. In another embodiment, the subject or subject group being treated in accordance with methods of the invention exhibits a baseline fasting plasma level (or mean thereof) of free EPA, expressed as a percentage of total free fatty acid, of not more than about 3%, not more than about 2.5%, not more than about 2%, not more than about 1.5%, not more than about 1%, not more than about 0.75%, not more than about 0.5%, not more than about 0.25%, not more than about 0.2% or not more than about 0.15%. In one such embodiment, free plasma EPA and/or total fatty acid levels are determined prior to initiating therapy.

**[0076]** In another embodiment, the subject or subject group being treated in accordance with methods of the invention exhibits a fasting baseline absolute plasma level of total fatty acid (or mean thereof) not greater than about 250 nmol/ml, not greater than about 200 nmol/ml, not greater than about 150 nmol/ml, not greater than about 100 nmol/ml, or not greater than about 50 nmol/ml.

**[0077]** In another embodiment, the subject or subject group being treated in accordance with methods of the invention exhibits a fasting baseline plasma, serum or red blood cell membrane EPA level not greater than about 70 µg/ml, not greater than about 60 µg/ml, not greater than about 50 µg/ml, not greater than about 40 µg/ml, not greater than about 30 µg/ml, or not greater than about 25 µg/ml.

**[0078]** In another embodiment, methods of the present invention comprise a step of measuring the subject’s (or subject group’s mean) baseline lipid profile prior to initiating therapy. In another embodiment, methods of the invention comprise the step of identifying a subject or subject group having one or more of the following: baseline non-HDL-C value of about 200 mg/dl to about 400 mg/dl, for example at least about 210 mg/dl, at least about 220 mg/dl, at least about 230 mg/dl, at least about 240 mg/dl, at least about 250 mg/dl, at least about 260 mg/dl, at least about 270 mg/dl, at least about 280 mg/dl, at least about 290 mg/dl, or at least about 300 mg/dl; baseline total cholesterol value of about 250 mg/dl to about 400 mg/dl, for example at least about 260 mg/dl, at least about 270 mg/dl, at least about 280 mg/dl or at least about 290 mg/dl; baseline vLDL-C value of about 140 mg/dl to about 200 mg/dl, for example at least about 150 mg/dl, at least about 160 mg/dl, at least about 170 mg/dl, at least about 180 mg/dl or at least about 190 mg/dl; baseline HDL-C value of about 10 to about 60 mg/dl, for example not more than about 40 mg/dl, not more than about 35 mg/dl, not more than about 30 mg/dl, not more than about 25 mg/dl, not more than about 20 mg/dl, or not more than about 15 mg/dl; and/or baseline LDL-C value of about 50 to about 300 mg/dl, for example not less than about 100 mg/dl, not less than about 90 mg/dl, not less than about 80 mg/dl,



not less than about 70 mg/dl, not less than about 60 mg/dl or not less than about 50 mg/dl.

[0079] In one embodiment, compositions of the invention are packaged in blister packs. In another embodiment, the blister packs comprise PCTFE (for example 50 $\mu$ ) laminated with water based adhesive to clear PVC (for example 1900 which are heat sealed to aluminum foil).

[0080] In a related embodiment, upon treatment in accordance with the present invention, for example over a period of about 1 to about 200 weeks, about 1 to about 100 weeks, about 1 to about 80 weeks, about 1 to about 50 weeks, about 1 to about 40 weeks, about 1 to about 20 weeks, about 1 to about 15 weeks, about 1 to about 12 weeks, about 1 to about 10 weeks, about 1 to about 5 weeks, about 1 to about 2 weeks or about 1 week, the subject or subject group exhibits one or more of the following outcomes:

- [0081] (a) reduced triglyceride levels compared to baseline or a placebo arm;
- [0082] (b) reduced Apo B levels compared to baseline or a placebo arm;
- [0083] (c) increased HDL-C levels compared to baseline or a placebo arm;
- [0084] (d) no increase in LDL-C levels compared to baseline or a placebo arm;
- [0085] (e) a reduction in LDL-C levels compared to baseline or a placebo arm;
- [0086] (f) a reduction in non-HDL-C levels compared to baseline or a placebo arm;
- [0087] (g) a reduction in vLDL levels compared to baseline or a placebo arm;
- [0088] (h) an increase in apo A-I levels compared to baseline or a placebo arm;
- [0089] (i) an increase in apo A-I/apo B ratio compared to baseline or a placebo arm;
- [0090] (j) a reduction in lipoprotein A levels compared to baseline or a placebo arm;
- [0091] (k) a reduction in LDL particle number compared to baseline or a placebo arm;
- [0092] (l) an increase in mean LDL size compared to baseline or a placebo arm;
- [0093] (m) a reduction in remnant-like particle cholesterol compared to baseline or a placebo arm;
- [0094] (n) a reduction in oxidized LDL compared to baseline or a placebo arm;
- [0095] (o) no change or a reduction in fasting plasma glucose (FPG) compared to baseline or a placebo arm;
- [0096] (p) a reduction in hemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>) compared to baseline or a placebo arm;
- [0097] (q) a reduction in homeostasis model insulin resistance compared to baseline or a placebo arm;
- [0098] (r) a reduction in lipoprotein associated phospholipase A2 compared to baseline or a placebo arm;
- [0099] (s) a reduction in intracellular adhesion molecule-1 compared to baseline or a placebo arm;
- [0100] (t) a reduction in interleukin-6 compared to baseline or a placebo arm;
- [0101] (u) a reduction in plasminogen activator inhibitor-1 compared to baseline or a placebo arm;
- [0102] (v) a reduction in high sensitivity C-reactive protein (hsCRP) compared to baseline or a placebo arm;
- [0103] (w) an increase in serum phospholipid EPA compared to baseline or a placebo arm;

[0104] (x) an increase in red blood cell membrane EPA compared to baseline or a placebo arm; and/or

[0105] (y) a reduction or increase in one or more of serum phospholipid and/or red blood cell content of docosahexaenoic acid (DHA), docosapentaenoic acid (DPA), arachidonic acid (AA), palmitic acid (PA), stearidonic acid (SA) or oleic acid (OA) compared to baseline or a placebo arm.

[0106] In one embodiment, methods of the present invention comprise measuring baseline levels of one or more markers set forth in (a)-(y) above prior to dosing the subject or subject group. In another embodiment, the methods comprise administering a composition as disclosed herein to the subject after baseline levels of one or more markers set forth in (a)-(y) are determined, and subsequently taking an additional measurement of said one or more markers.

[0107] In another embodiment, upon treatment with a composition of the present invention, for example over a period of about 1 to about 200 weeks, about 1 to about 100 weeks, about 1 to about 80 weeks, about 1 to about 50 weeks, about 1 to about 40 weeks, about 1 to about 20 weeks, about 1 to about 15 weeks, about 1 to about 12 weeks, about 1 to about 10 weeks, about 1 to about 5 weeks, about 1 to about 2 weeks or about 1 week, the subject or subject group exhibits any 2 or more of, any 3 or more of, any 4 or more of, any 5 or more of, any 6 or more of, any 7 or more of, any 8 or more of, any 9 or more of, any 10 or more of, any 11 or more of, any 12 or more of, any 13 or more of, any 14 or more of, any 15 or more of, any 16 or more of, any 17 or more of, any 18 or more of, any 19 or more of, any 20 or more of, any 21 or more of, any 22 or more of, any 23 or more, any 24 or more, or all 25 of outcomes (a)-(y) described immediately above.

[0108] In another embodiment, upon treatment with a composition of the present invention, the subject or subject group exhibits one or more of the following outcomes:

- [0109] (a) a reduction in triglyceride level of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) as compared to baseline or a placebo arm;
- [0110] (b) a less than 30% increase, less than 20% increase, less than 10% increase, less than 5% increase or no increase in non-HDL-C levels or a reduction in non-HDL-C levels of at least about 1%, at least about 3%, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) as compared to baseline or a placebo arm;
- [0111] (c) substantially no change, no change or an increase in HDL-C levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) as compared to baseline or a placebo arm;
- [0112] (d) a less than 60% increase, less than 50% increase, less than 40% increase, less than 30% increase, less than 20% increase, less than 10%



- increase, less than 5% increase or no increase in LDL-C levels or a reduction in LDL-C levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 55% or at least about 75% (actual % change or median % change) as compared to baseline or a placebo arm;
- [0113] (e) a decrease in Apo B levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) as compared to baseline or a placebo arm;
- [0114] (f) a reduction in vLDL levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0115] (g) an increase in apo A-I levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0116] (h) an increase in apo A-I/apo B ratio of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0117] (i) a reduction in lipoprotein(a) levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0118] (j) a reduction in mean LDL particle number of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0119] (k) an increase in mean LDL particle size of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0120] (l) a reduction in remnant-like particle cholesterol of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0121] (m) a reduction in oxidized LDL of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0122] (n) substantially no change, no change or a reduction in fasting plasma glucose (FPG) of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0123] (o) substantially no change, no change or a reduction in hemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>) of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, or at least about 50% (actual % change or median % change) compared to baseline or a placebo arm;
- [0124] (p) a reduction in homeostasis model index insulin resistance of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0125] (q) a reduction in lipoprotein associated phospholipase A2 of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0126] (r) a reduction in intracellular adhesion molecule-1 of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0127] (s) a reduction in interleukin-6 of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;
- [0128] (t) a reduction in plasminogen activator inhibitor-1 of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline;
- [0129] (u) a reduction in high sensitivity C-reactive protein (hsCRP) of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or a placebo arm;



[0130] (v) an increase in serum, plasma and/or RBC EPA of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 100%, at least about 200% or at least about 400% (actual % change or median % change) compared to baseline or a placebo arm;

[0131] (w) an increase in serum phospholipid and/or red blood cell membrane EPA of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 100%, at least about 200%, or at least about 400% (actual % change or median % change) compared to baseline or a placebo arm;

[0132] (x) a reduction or increase in one or more of serum phospholipid and/or red blood cell DHA, DPA, AA, PA and/or OA of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) compared to baseline or a placebo arm; and/or

[0133] (y) a reduction in total cholesterol of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) compared to baseline or a placebo arm.

[0134] In one embodiment, methods of the present invention comprise measuring baseline levels of one or more markers set forth in (a)-(y) prior to dosing the subject or subject group. In another embodiment, the methods comprise administering a composition as disclosed herein to the subject after baseline levels of one or more markers set forth in (a)-(y) are determined, and subsequently taking a second measurement of the one or more markers as measured at baseline for comparison thereto.

[0135] In another embodiment, upon treatment with a composition of the present invention, for example over a period of about 1 to about 200 weeks, about 1 to about 100 weeks, about 1 to about 80 weeks, about 1 to about 50 weeks, about 1 to about 40 weeks, about 1 to about 20 weeks, about 1 to about 15 weeks, about 1 to about 12 weeks, about 1 to about 10 weeks, about 1 to about 5 weeks, about 1 to about 2 weeks or about 1 week, the subject or subject group exhibits any 2 or more of, any 3 or more of, any 4 or more of, any 5 or more of, any 6 or more of, any 7 or more of, any 8 or more of, any 9 or more of, any 10 or more of, any 11 or more of, any 12 or more of, any 13 or more of, any 14 or more of, any 15 or more of, any 16 or more of, any 17 or more of, any 18 or more of, any 19 or more of, any 20 or more of, any 21 or more of, any 22 or more of, any 23 or more of, any 24 or more of, or all 25 of outcomes (a)-(y) described immediately above.

[0136] Parameters (a)-(y) can be measured in accordance with any clinically acceptable methodology. For example, triglycerides, total cholesterol, HDL-C and fasting blood sugar can be sample from serum and analyzed using standard photometry techniques. VLDL-TG, LDL-C and VLDL-C can be calculated or determined using serum

lipoprotein fractionation by preparative ultracentrifugation and subsequent quantitative analysis by refractometry or by analytic ultracentrifugal methodology. Apo A1, Apo B and hsCRP can be determined from serum using standard nephelometry techniques. Lipoprotein (a) can be determined from serum using standard turbidimetric immunoassay techniques. LDL particle number and particle size can be determined using nuclear magnetic resonance (NMR) spectrometry. Remnants lipoproteins and LDL-phospholipase A2 can be determined from EDTA plasma or serum and serum, respectively, using enzymatic immunoseparation techniques. Oxidized LDL, intercellular adhesion molecule-1 and interleukin-2 levels can be determined from serum using standard enzyme immunoassay techniques. These techniques are described in detail in standard textbooks, for example Tietz Fundamentals of Clinical Chemistry, 6th Ed. (Burtis, Ashwood and Bortor Eds.), WB Saunders Company.

[0137] In one embodiment, subjects fast for up to 12 hours prior to blood sample collection, for example about 10 hours.

[0138] In another embodiment, the present invention provides a method of treating or preventing primary hypercholesterolemia and/or mixed dyslipidemia (Fredrickson Types IIa and IIb) in a patient in need thereof, comprising administering to the patient one or more compositions as disclosed herein. In a related embodiment, the present invention provides a method of reducing triglyceride levels in a subject or subjects when treatment with a statin or niacin extended-release monotherapy is considered inadequate (Frederickson type IV hyperlipidemia).

[0139] In another embodiment, the present invention provides a method of treating or preventing risk of recurrent nonfatal myocardial infarction in a patient with a history of myocardial infarction, comprising administering to the patient one or more compositions as disclosed herein.

[0140] In another embodiment, the present invention provides a method of slowing progression of or promoting regression of atherosclerotic disease in a patient in need thereof, comprising administering to a subject in need thereof one or more compositions as disclosed herein.

[0141] In another embodiment, the present invention provides a method of treating or preventing very high serum triglyceride levels (e.g. Types IV and V hyperlipidemia) in a patient in need thereof, comprising administering to the patient one or more compositions as disclosed herein.

[0142] In another embodiment, the present invention provides a method of treating subjects having very high serum triglyceride levels (e.g. greater than 1000 mg/dl or greater than 2000 mg/dl) and that are at risk of developing pancreatitis, comprising administering to the patient one or more compositions as disclosed herein.

[0143] In one embodiment, a composition of the invention is administered to a subject in an amount sufficient to provide a daily dose of eicosapentaenoic acid of about 1 mg to about 10,000 mg, 25 about 5000 mg, about 50 to about 3000 mg, about 75 mg to about 2500 mg, or about 100 mg to about 1000 mg, for example about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, about 300 mg, about 325 mg, about 350 mg, about 375 mg, about 400 mg, about 425 mg, about 450 mg, about 475 mg, about 500 mg, about 525 mg, about 550 mg, about 575 mg, about 600 mg, about 625 mg, about 650 mg, about 675 mg, about 700 mg, about 725 mg, about 750 mg, about 775 mg, about 800 mg,



about 825 mg, about 850 mg, about 875 mg, about 900 mg, about 925 mg, about 950 mg, about 975 mg, about 1000 mg, about 1025 mg, about 1050 mg, about 1075 mg, about 1100 mg, about 1025 mg, about 1050 mg, about 1075 mg, about 1200 mg, about 1225 mg, about 1250 mg, about 1275 mg, about 1300 mg, about 1325 mg, about 1350 mg, about 1375 mg, about 1400 mg, about 1425 mg, about 1450 mg, about 1475 mg, about 1500 mg, about 1525 mg, about 1550 mg, about 1575 mg, about 1600 mg, about 1625 mg, about 1650 mg, about 1675 mg, about 1700 mg, about 1725 mg, about 1750 mg, about 1775 mg, about 1800 mg, about 1825 mg, about 1850 mg, about 1875 mg, about 1900 mg, about 1925 mg, about 1950 mg, about 1975 mg, about 2000 mg, about 2025 mg, about 2050 mg, about 2075 mg, about 2100 mg, about 2125 mg, about 2150 mg, about 2175 mg, about 2200 mg, about 2225 mg, about 2250 mg, about 2275 mg, about 2300 mg, about 2325 mg, about 2350 mg, about 2375 mg, about 2400 mg, about 2425 mg, about 2450 mg, about 2475 mg or about 2500 mg.

[0144] In another embodiment, any of the methods disclosed herein are used in treatment or prevention of a subject or subjects that consume a traditional Western diet. In one embodiment, the methods of the invention include a step of identifying a subject as a Western diet consumer or prudent diet consumer and then treating the subject if the subject is deemed a Western diet consumer. The term “Western diet” herein refers generally to a typical diet consisting of, by percentage of total calories, about 45% to about 50% carbohydrate, about 35% to about 40% fat, and about 10% to about 15% protein. A Western diet may alternately or additionally be characterized by relatively high intakes of red and processed meats, sweets, refined grains, and deserts, for example more than 50%, more than 60% or more or 70% of total calories come from these sources.

EXAMPLES

[0145] The following examples are for illustrative purposes only and should not be construed as limiting the invention in any manner.

Example 1

[0146] A Test Composition (TC) was prepared comprising ultra-pure Ethyl-EPA (>96% E-EPA, ~3% related fatty acid substances (no DHA), and ~0.2% alpha tocopherol) filled into soft gelatin capsule shells (~500 mg fill weight per capsule) prepared from a gel comprising gelatin (~44%), glycerol (~7%), sorbitol solution (~17%), maltitol solution, gelatin and purified water. A Comparative Composition (CC) was made comprising the same fill as the Test Composition but filled into Type IIa Capsules made from a gel comprising of glycerol (~20%), gelatin (43.4%) and water (~36.6%).

[0147] Test Compositions and Comparative Compositions were then placed in polybags which were sealed and stored at either 25° C./60% RH or 30° C./65% RH for a period of 1, 3, or 6 months. At the end of storage, capsules were opened and peroxide value of the fill material was analyzed. Results are shown in Table 1 (average of capsules from three different batches).

TABLE 1

Peroxide Values (meq/mg) Upon Storage.				
Composition	Baseline	1 Month	3 Month	6 Month
Storage at 25° C./60% RH				
TC	1.6	—	3.2	3.4
CC	1.9	—	3.4	9.6
Storage at 30° C./65% RH				
TC	1.6	2.0	3.6	4.8
CC	1.8	1.9	3.5	12.5

[0148] As is seen in Table 1, the Test Composition fill material exhibited much lower peroxide values after 6 months of storage under both sets of storage conditions. No significant differences were observed between the Test Composition and Comparative Composition fill material in terms of potency of EPA-E and related substances throughout the duration of the study.

Example 2

[0149] Test Compositions and Comparative Compositions of Example 1 were prepared and packaged in blister packaging (50μ PCTFE laminated with water based adhesive to 190μ clear PVC and heat sealed to aluminum foil). Packaged Test Compositions and Comparative Compositions were then stored at either 25° C./60% RH or 40° C./70% RH for a period of 1, 3, 6, 12 or 36 months. At the end of storage, capsules were opened and the peroxide values of the fill contents analyzed as shown in Table 2 (average of the three batches).

TABLE 2

Peroxide Values (meq/mg) Upon Storage.						
	Baseline	1 Mo.	3 Mo.	6 Mo.	9 Mo.	12 Mo.
Storage at 25° C./60% RH						
TC	2.5	—	1.1	2.1	2.2	5.4
CC	2.6	—	5.1	8.3	9.7	11.1
Storage at 40° C./75% RH						
TC	2.5	2.1	3.2	4.9	—	—
CC	2.6	3.4	10.6	18.8	—	—

[0150] As is seen in Table 2, the Test Composition exhibited much lower peroxide values after 3, 6, 9 and 12 months of storage at 25° C./60% RH and after 1, 3 and 6 months of storage at 40° C./75% RH as compared to the Comparative Compositions.

[0151] At 40° C., the Test Composition showed an average decrease in E-EPA potency of 0.30% per month whereas the Comparative Compositions showed an average decrease in E-EPA potency of 0.44% per month. However, similar results were not obtained with the same batches in Example 1 (not stored in blister packages). Additionally, the related substances measurements did not show any concomitant increase suggesting normal analytic variation may be responsible.

[0152] When the peroxide values were forced to linear trendlines, average slope values between the Test Compositions in Experiment 1 (no blister packaging) and Experi-



ment 2 (blister packaging) were similar indicating that the packaging is likely not responsible for prevention of oxidation.

TABLE 3

Peroxide Value: Linear Slope Comparison Between Example 1 and Example 2				
Storage Conditions	Test Composition Slope (meq/kg/mo.)		Comparative Composition Slope (meq/kg/mo.)	
	Example 1	Example 2	Example 1	Example 2
25° C./60% RH	0.33	0.35	1.45	1.03
40° C./75% RH	0.56	0.66	1.81	3.00

Example 3

[0153] A dissolution test was performed on capsules of Example 1 containing 500 mg E-EPA using the Rotating Dialysis Cell method set forth in Yamazaki et al., *Dissolution tests by RDC method for soft gelatin capsules containing ethyl icosapentate*, Pharmaceutical Technology Japan, 15: 595-603 (1999). Conditions were as set forth below:

- [0154] RDC Cell: PharmaTest
- [0155] Paddle speed: 100 rpm
- [0156] Temperature: 37° C.
- [0157] Filter: Millipore hydrophobic filter sheets
- [0158] Inner media: JP pH 1.2 disintegration media
- [0159] Outer Media: Absolute Ethanol
- [0160] Samples: 5 ml taken at 10, 20, 30, 40, 60, 100 and 120 minutes

[0161] The samples were analyzed against a reference standard prepared in ethanol at 0.5 mg/ml, the amount of product dissolved at each time point was then calculated. A good dissolution profile was obtained with a Q85 of approximately 60 minutes and a profile very similar to that generated by Yamazaki (JP data; succinated gelatin capsules). Dissolution profile of the inventive capsule composition was also evaluated by the paddle method in media containing buffer, SDS and IPA (100 rpm paddle speed, 1000 ml, 37° C.). Samples were removed at intervals and analyzed against a standard solution (9.5 ml/ml in methanol) by HPCL. All data are shown in FIG. 1.

Example 4

[0162] Bioavailability data were obtained for a capsule shell according to Example 1 containing 500 mg E-EPA (AMR101) and were compared against reported by Yamazaki data for 300 mg Epadel capsules (succinated gelatin; Comparitor 1 and Comparitor 2). Tmax data are shown in Table 4 together with dissolution percentage at 60 min. Full bioavailability profiles for EPA succinated capsules and AMR101 capsules are shown in FIGS. 2 and 3, respectively.

TABLE 4

Dissolution and Tmax.		
	Dissolution at 60 min (%)	T <sub>max</sub> (hrs)
Comparitor 1 <sup>1</sup>	77	6
Comparitor 2 <sup>2</sup>	75	6
AMR101 <sup>3</sup>	87	5

<sup>1</sup>Capsule shell = 220 mg; contents = 323 mg.  
<sup>2</sup>Capsule shell = 134 mg; contents = 327 mg.  
<sup>3</sup>Mean of three batches using RDC and pH 1.2.

[0163] As can be seen from Table 4, AMR101 exhibited greater E-EPA dissolution by 60 minutes, and had a shorter T<sub>max</sub> than was reported for Epadel present in succinated gelatin capsules.

- 1-10. (canceled)
- 11. A pharmaceutical composition comprising ethyl eicosapentaenoate (EPA-E), lecithin, and one or more surfactants, wherein the EPA-E is present in an amount of at least about 90% by weight of all omega-3 fatty acids in the composition.
- 12. The pharmaceutical composition of claim 11, wherein the EPA-E is present in an amount of at least about 96 wt. % of all omega-3 fatty acids in the composition.
- 13. The pharmaceutical composition of claim 11, wherein the pharmaceutical composition comprises the composition contains not more than 0.6% of any individual fatty acid other than EPA-E.
- 14. The pharmaceutical composition of claim 11, wherein the pharmaceutical composition is free of docosahexaenoic acid.
- 15. The pharmaceutical composition of claim 11, wherein the composition comprises two surfactants.
- 16. The pharmaceutical composition of claim 11, wherein the pharmaceutical composition is encapsulated in a capsule shell.
- 17. The pharmaceutical composition of claim 11, wherein the EPA-E is present in the pharmaceutical composition in an amount of from about 100 mg to about 1000 mg.
- 18. The pharmaceutical composition of claim 17, wherein the EPA-E is present in the pharmaceutical composition in an amount of about 800 mg.
- 19. A method of lowering triglycerides in a subject with fasting triglycerides of from about 500 mg/dl to about 2500 mg/dL, comprising orally administering to the subject daily a pharmaceutical composition comprising ethyl eicosapentaenoate (EPA-E), lecithin, and one or more surfactants, wherein the EPA-E is present in an amount of at least about 90% by weight of all omega-3 fatty acids in the composition for a period effective to reduce triglycerides in the subject without increasing LDL-C by more than 20%.
- 20. The method of claim 19, wherein the subject has fasting serum triglycerides of at least 500 mg/dL.
- 21. The method of claim 19, wherein the subject has fasting serum triglycerides of at least 700 mg/dL.
- 22. The method of claim 19, comprising administering to the subject the pharmaceutical composition daily for a period effective to reduce triglycerides by at least 10% in the subject.
- 23. The method of claim 19, comprising administering to the subject the pharmaceutical composition daily for a period effective to reduce triglycerides by at least 20% in the subject.



**24.** The method of claim **19**, comprising administering to the subject the pharmaceutical composition daily for a period effective to reduce triglycerides without increasing LDL-C by more than 10%.

**25.** The method of claim **19**, comprising administering to the subject the pharmaceutical composition daily for a period effective to reduce non-HDL-C by 5% in the subject.

**26.** The method of claim **19**, comprising administering to the subject the pharmaceutical composition daily for a period effective to increase HDL-C in the subject.

**27.** The method of claim **19**, wherein the EPA-E is present in an amount of at least about 96 wt. % of all omega-3 fatty acids in the composition.

**28.** The method of claim **19**, wherein the pharmaceutical composition comprises the composition contains not more than 0.6% of any individual fatty acid other than EPA-E.

**29.** The method of claim **19**, wherein the EPA-E is present in the pharmaceutical composition in an amount of from about 100 mg to about 1000 mg.

**30.** The method of claim **19**, wherein the EPA-E is present in the pharmaceutical composition in an amount of about 800 mg.

\* \* \* \* \*